European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Supporting documentation for the conservation status assessment for the habitat:

H2150 - Atlantic decalcified fixed dunes (Calluno-Ulicetea)

SCOTLAND

IMPORTANT NOTE - PLEASE READ

- The information in this document is a country-level contribution to the UK Report on the conservation status of this habitat, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- The 2019 Article 17 UK Approach document provides details on how this supporting information was used to produce the UK Report.
- The UK Report on the conservation status of this habitat is provided in a separate document.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Explanatory notes (where provided) by the country are included at the end. These provide an audit trail of relevant supporting information.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was only relevant at UK-level (sections 10 Future prospects and 11 Conclusions).
- For technical reasons, the country-level future trends for Range, Area covered by habitat and Structure and functions are only available in a separate spreadsheet that contains all the country-level supporting information.
- The country-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

NATIONAL LEVEL

1. General information

1.1 Member State	UK (Scotland information only)
1.2 Habitat code	2150 - Atlantic decalcified fixed dunes (Calluno-Ulicetea)

2. Maps

2.1 Year or period	1987-2006
2.3 Distribution man	Ves

2.3 Distribution map Method used Complete survey or a statistically robust estimate

2.4 Additional maps

BIOGEOGRAPHICAL LEVEL

3. Biogeographical and marine regions

3.1 Biogeographical or marine region where the habitat occurs

3.2 Sources of information

Atlantic (ATL)

https://www.environment.gov.scot/our-environment/habitats-and-species/habitat-map-of-scotland/

The Sand Dune Vegetation Survey of Scotland, Tom Dargie 1994-2000 National Report

The Sand Dune Vegetation Survey of Scotland 2012 SNH Natural Spaces dataset Janine M Morris, Site Condition Monitoring of Coastal Habitats. (National Contract, Year 2009-2010) and Site Condition Monitoring of Coastal Habitats (National Contract, Year 2010-2011). Contract No: 25639

SNH Site Condition Monitoring results Cycle 3 (from 1 April 2012): see Scotland's environment website. [From the website Detailed tab, select Coastal features by clicking the Feature filter on the left of the screen, then Feature Category= Coast. Data can be exported to spreadsheet by right clicking the table at the bottom of the screen, then Export, then Export Table. Cycle 3 assessments can be seen by filtering the spreadsheet on the 'LatestAssessedSCMcycle' column].

http://jncc.defra.gov.uk/pdf/Article17Consult_20131010/H2150_SCOTLAND.pdf

4. Range

- 4.1 Surface area (in km²)
- 4.2 Short-term trend Period
- 4.3 Short-term trend Direction
- 4.4 Short-term trend Magnitude
 4.5 Short-term trend Method used
- - ·
- 4.6 Long-term trend Period
- 4.7 Long-term trend Direction
- 4.8 Long-term trend Magnitude
- 4.9 Long-term trend Method used
- 4.10 Favourable reference range

- Stable (0)
- a) Minimum

b) Maximum

- a) Minimum
- b) Maximum
- a) Area (km²)
- b) Operator
- c) Unknown No
- d) Method

Improved knowledge/more accurate data

The change is mainly due to: Improved knowledge/more accurate data

4.11 Change and reason for change in surface area of range

4.12 Additional information

Final data has become available from translation of the Sand Dune Survey of Scotland to Annex I habitats as well as some other NVC surveys and this has given a complete picture of the distribution of this habitat in Scotland. The changes from previous reported distribution are not great however.

5. Area covered by habitat

iod

1987-2006

5.2 Surface area (in km²)

a) Minimum

b) Maximum

c) Best single 11.73

value

5.3 Type of estimate

5.4 Surface area Method used

5.5 Short-term trend Period

5.6 Short-term trend Direction

5.7 Short-term trend Magnitude

95% confidence interval

Complete survey or a statistically robust estimate

2005-2016

Stable (0)

a) Minimum

b) Maximum

Based mainly on extrapolation from a limited amount of data

c) Confidence

interval

5.8 Short-term trend Method used

5.9 Long-term trend Period

5.10 Long-term trend Direction

5.11 Long-term trend Magnitude

a) Minimum

b) Maximum

c) Confidence

interval

5.12 Long-term trend Method used

5.13 Favourable reference area

a) Area (km²)

b) Operator

c) Unknown No

d) Method

5.14 Change and reason for change in surface area of range

Improved knowledge/more accurate data

The change is mainly due to:

Improved knowledge/more accurate data

5.15 Additional information

The results of translating the Sand Dune Survey of Scotland from NVC to Annex I habitats has become available through HabMoS. This has given a more accurate data on extent of this habitat, which is greater than previously reported. Other NVC survey data has also become available translated rom NVC through HabMoS. There are still some issues with the data from HaBMoS (some overlapping polygons) but this is our best current data.

6. Structure and functions

6.1 Condition of habitat

a) Area in good condition

Minimum 2.9

Maximum 2.9

(km²)

b) Area in not-good

not known (km²)

Minimum 5.29

Maximum 5.29

condition (km²)

c) Area where condition is

Minimum 3.55

Based mainly on extrapolation from a limited amount of data

Maximum 3.55

6.2 Condition of habitat Method used

6.3 Short-term trend of habitat area in good condition Period

6.4 Short-term trend of habitat area in good condition Direction

Decreasing (-)

2001-2016

6.5 Short-term trend of habitat area in good condition Method used

Based mainly on extrapolation from a limited amount of data

6.6 Typical species

Has the list of typical species changed in comparison to the previous No reporting period?

6.7 Typical species Method used6.8 Additional information

7. Main pressures and threats

7.1 Characterisation of pressures/threats

Pressure	Ranking
Mixed source air pollution, air-borne pollutants (J03)	Н
Extensive grazing or undergrazing by livestock (A10)	Н
Intensive grazing or overgrazing by livestock (A09)	M
Problematic native species (I04)	Н
Natural succession resulting in species composition change (other than by direct changes of agricultural or forestry practices) (LO2)	M
Threat	Ranking
Mixed source air pollution, air-borne pollutants (J03)	Н
Extensive grazing or undergrazing by livestock (A10)	Н
Intensive grazing or overgrazing by livestock (A09)	M
Problematic native species (I04)	Н

M

7.2 Sources of information

practices) (L02)

7.3 Additional information

8. Conservation measures

Natural succession resulting in species composition change

(other than by direct changes of agricultural or forestry

8.1 Status of measures	a) Are measures needed?	Yes
	b) Indicate the status of measures	Measures identified and taken
8.2 Main purpose of the measures taken	Maintain the current range, populati	on and/or habitat for the species
8.3 Location of the measures taken	Both inside and outside Natura 2000	
8.4 Response to the measures	Medium-term results (within the next two reporting periods, 2019-2030)	
8.5 List of main conservation measures		

Management of habitats (others than agriculture and forest) to slow, stop or reverse natural processes (CL01)

Management of problematic native species (CI05)

8.6 Additional information

Although there are concerns about negative trends in the structure and function of the habitat, remedies have been put in place through management plans on military sites and SSSI agreements on private land. With these recovery actions in place at a number of sites the situation should begin to improve

9. Future prospects

9.1 Future prospects of parameters

- a) Range
- b) Area
- c) Structure and functions

9.2 Additional information

No evidence to suggest that problems with scrub encroachment, self sown trees and invasion by Chamaerion angustifolium are not ongoing. Nitrogen deposition will likely have a high impact

10. Conclusions

10.1. Range

10.2. Area

10.3. Specific structure and functions

(incl. typical species)

10.4. Future prospects

10.5 Overall assessment of

Conservation Status

10.6 Overall trend in Conservation Status

10.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

10.8 Additional information

11. Natura 2000 (pSCIs, SCIs, SACs) coverage for Annex I habitat types

11.1 Surface area of the habitat type inside the pSCIs, SCIs and SACs network (in km² in biogeographical/marine region)

a) Minimum

b) Maximum

c) Best single value 3.51

11.2 Type of estimate

11.3 Surface area of the habitat type inside the network Method used

11.4 Short-term trend of habitat area in good condition within the network Direction

11.5 Short-term trend of habitat area in good condition within network Method used

11.6 Additional information

c) best siligle

Best estimate

Complete survey or a statistically robust estimate

Stable (0)

Based mainly on extrapolation from a limited amount of data

12. Complementary information

12.1 Justification of % thresholds for trends

12.2 Other relevant information

Distribution Map




Figure 1: UK distribution map for H2150 - Atlantic decalcified fixed dunes (*Calluno-Ulicetea*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available habitat records which are considered to be representative of the distribution within the current reporting period. For further details see the 2019 Article17 UK Approach document.

Range Map




Figure 2: UK range map for H2150 - Atlantic decalcified fixed dunes (*Calluno-Ulicetea*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by applying a bespoke range mapping tool for Article 17 reporting (produced by JNCC) to the 10km grid square distribution map presented in Figure 1. The alpha value for this habitat was 25km. For further details see the 2019 Article 17 UK Approach document.