# European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

# Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Supporting documentation for the conservation status assessment for the habitat:

H3130 - Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or of the *Isoëto-Nanojuncetea* 

NORTHERN IRELAND

#### **IMPORTANT NOTE - PLEASE READ**

- The information in this document is a country-level contribution to the UK Report on the conservation status of this habitat, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- The 2019 Article 17 UK Approach document provides details on how this supporting information was used to produce the UK Report.
- The UK Report on the conservation status of this habitat is provided in a separate document.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Explanatory notes (where provided) by the country are included at the end. These provide an audit trail of relevant supporting information.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was only relevant at UK-level (sections 10 Future prospects and 11 Conclusions).
- For technical reasons, the country-level future trends for Range, Area covered by habitat and Structure and functions are only available in a separate spreadsheet that contains all the country-level supporting information.
- The country-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

#### **NATIONAL LEVEL**

#### 1. General information

1.1 Member State	UK (Northern Ireland information only)
1.2 Habitat code	3130 - Oligotrophic to mesotrophic standing waters with vegetation of the Lit

#### 2. Maps

2.1 Year or period	2007-2017
2.3 Distribution map	Yes

2.3 Distribution map Method used Complete survey or a statistically robust estimate

2.4 Additional maps

#### BIOGEOGRAPHICAL LEVEL

#### 3. Biogeographical and marine regions

3.1 Biogeographical or marine region where the habitat occurs

3.2 Sources of information

#### Atlantic (ATL)

JOINT NATURE CONSERVATION COMMITTEE 2005. Common Standards Monitoring (CSM). Joint Nature Conservation Committee, Peterborough. www.jncc.gov.uk/page-2217

PALMER, M.A., BELL, S.L. & BUTTERFIELD, I. 1992. A botanical classification of standing waters in Britain: applications for conservation and monitoring. Aquatic Conservation: Marine and Freshwater Ecosystems 2: 125 - 143.

PALMER, M.A. & ROY, D.B. 2001a. A method for estimating the extent of standing fresh waters of different trophic states in Great Britain. Aquatic Conservation: Marine and Freshwater Ecosystems, 11, 199-216.

http://www3.interscience.wiley.com/cgi-bin/abstract/83502064/START PALMER, M.A. & ROY, D.B. 2001b.

Second Report by the United Kingdom under Article 17 on the implementation of the Directive from January 2001 to December 2006

WILLIAMS, J.M. (ed.) 2006. Common Standards Monitoring for Designated Sites: First Six Year Report. Joint Nature Conservation Committee, Peterborough. http://www.jncc.gov.uk/page-3520

WOLFE-MURPHY, S.A., LAWRIE, E.W., SMITH, S.J. & GIBSON, C.E. 1992. Northern Ireland Lakes Survey. Unpublished report to Northern Ireland Department of Environment, Belfast.

ENSIS Ltd (Environmental Science Services) Goldsmith, B., Davidson, T. A., Burgess, A., Hughes, M., Madgwick, G., Rawcliffe, R., Rippey, B. & Tyler, J., December 2008 Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland. Final Report to Northern Ireland Environment Agency. POND CONSERVATION Helen Keeble, Penny Williams, Jeremy Biggs & Neil Reid, 2009, Important Areas for Ponds (IAPs) and other small waterbodies in Northern Ireland, NIEA Research and Development Series 11/5.

MC ELARNEY, Y.R. FOY, R.H. PARK, R. ANDERSON, N.J. PLA-RABES, S.RASMUSSEN, P.O'DEA, P.ENGSTOM, D.R. & MCGOWAN, S. 2009. A framework for the management of forest inpacts on upland lakes INTERREG - Project 20274 MC ELARNEY, Y.R. FOY, R. ANDERSON, S. & RASMUSSEN, P. 2010. Response of aquatic macrophytes in Northern Ireland softwater lakes to forestry management; eutrophication and dissolved organic carbon. Aquatic Botany AQBOT - 2335. www.elsevier.com/locate/aquabot. Northern Ireland Environment Agency unpublished survey and monitoring data 2000-2012.

Third Report by the United Kingdom under Article 17 on the implementation of the Directive from January 2007 to December 2012. Printed on 06/11/2013 Page 5

JNCC Common Standards Monitoring Guidance for Freshwater Habitats and Species, Rivers and Lakes guidance updated September 2016 and March 2015 respectively, ISSN 1743-8160 jnccdefra.gov.uk/page - 2231

Goldsmith, B., Dowman, S., Goodrich, S., Shilland, E. & Shilland, J. (2015) DOE NIEA Standing Fresh Water Monitoring of Special Areas of Conservation (SAC) and Areas Special Scientific Interest (ASSI) (Second Round). P\_12324. ECRC (Environmental Change Research Centre) Research Report 168.

Air Pollution Information System (APIS). Http://.apis.ac.uk

Committee on Climate Change. 2017. UK Climate Change Risk Assessment (CCRAR) Evidence Report, Summary for Northern Ireland. Https://www.theecc.gov.uk

NORTHERN IRELAND ENVIRONMENT AGENCY. Nov 2014. Operational Policy Guidance Note on Alignment of Environmental Standards for Assessment under Habitats Directive and Water Framework Directive. DO1/13/483427

#### 4. Range

- 4.1 Surface area (in km²)
- 4.2 Short-term trend Period
- 4.3 Short-term trend Direction
- 4.4 Short-term trend Magnitude
- 4.5 Short-term trend Method used
- 4.6 Long-term trend Period
- 4.7 Long-term trend Direction
- 4.8 Long-term trend Magnitude
- 4.9 Long-term trend Method used
- 4.10 Favourable reference range
- 4.11 Change and reason for change

Stable (0)

a) Minimum

b) Maximum

1994-2018

Stable (0)

a) Minimum

b) Maximum

Complete survey or a statistically robust estimate

- a) Area (km²)
- b) Operator
- c) Unknown No
- d) Method

No change

The change is mainly due to:

#### 4.12 Additional information

in surface area of range

#### 5. Area covered by habitat

5.1 Year or period

5.2 Surface area (in km²)

2013-2017

a) Minimum

b) Maximum

c) Best single 23.59

value

5.3 Type of estimate

5.4 Surface area Method used

5.5 Short-term trend Period

5.6 Short-term trend Direction

5.7 Short-term trend Magnitude

95% confidence interval

Complete survey or a statistically robust estimate

2007-2018 Stable (0)

. . . . .

a) Minimum b) Maximum

c) Confidence interval

5.8 Short-term trend Method used

5.9 Long-term trend Period

Complete survey or a statistically robust estimate

1994-2018

Timex D <sub>j</sub>		
Stable (0)		
a) Minimum	b) Maximum	c) Confidence interval
Complete survey	or a statistically robust estimate	
<ul><li>a) Area (km²)</li><li>b) Operator</li><li>c) Unknown</li><li>d) Method</li></ul>	No	
No change		
The change is ma	ainly due to:	
	Stable (0) a) Minimum  Complete survey a) Area (km²) b) Operator c) Unknown d) Method No change	Stable (0) a) Minimum b) Maximum  Complete survey or a statistically robust estimate a) Area (km²) b) Operator c) Unknown No d) Method

5.15 Additional information

#### 6. Structure and functions

6.1 Condition of habitat	a) Area in good condition (km²)	Minimum 3	Maximum 3
	b) Area in not-good condition (km²)	Minimum 18.27	Maximum 18.27
	c) Area where condition is not known (km²)	Minimum 2.32	Maximum 2.32
6.2 Condition of habitat Method used	Based mainly on extrapolat	ion from a limited amo	ount of data
6.3 Short-term trend of habitat area in good condition Period	2007-2018		
6.4 Short-term trend of habitat area in good condition Direction	Decreasing (-)		
6.5 Short-term trend of habitat area	Based mainly on extrapolat	ion from a limited amo	ount of data
in good condition Method used	Has the list of typical specie	s changed in comparis	on to the previous No
6.6 Typical species	reporting period?		on to the previous No
6.7 Typical species Method used			
6.8 Additional information			

#### 7. Main pressures and threats

#### 7.1 Characterisation of pressures/threats

Pressure	Ranking
Intensive grazing or overgrazing by livestock (A09)	M
Extraction of minerals (e.g. rock, metal ores, gravel, sand, shell) (CO1)	М
Peat extraction (C05)	M
Mixed source pollution to surface and ground waters (limnic and terrestrial) (J01)	Н
Mixed source air pollution, air-borne pollutants (J03)	M
Temperature changes (e.g. rise of temperature & extremes) due to climate change (N01)	М
Increases or changes in precipitation due to climate change (N03)	М

Modification of hydrological flow (K04)	M
Abstraction from groundwater, surface water or mixed water (K01)	M
Other invasive alien species (other then species of Union concern) (IO2)	Н
Threat	Ranking
Intensive grazing or overgrazing by livestock (A09)	M
Extraction of minerals (e.g. rock, metal ores, gravel, sand, shell) (CO1)	M
Peat extraction (C05)	M
Mixed source pollution to surface and ground waters (limnic and terrestrial) (J01)	Н
Mixed source air pollution, air-borne pollutants (J03)	M
Temperature changes (e.g. rise of temperature & extremes) due to climate change (N01)	Н
Increases or changes in precipitation due to climate change (N03)	Н
Modification of hydrological flow (K04)	M
Abstraction from groundwater, surface water or mixed water (K01)	M
Other invasive alien species (other then species of Union concern) (IO2)	Н

7.2 Sources of information

7.3 Additional information

#### 8. Conservation measures

8.1 Status of measures	a) Are measures needed?	Yes
	b) Indicate the status of measures	Measures identified and taken
8.2 Main purpose of the measures taken	Restore the habitat of the species (re	elated to 'Habitat for the species')
8.3 Location of the measures taken	Both inside and outside Natura 2000	
8.4 Response to the measures	Medium-term results (within the nex	xt two reporting periods, 2019-2030)
8.5 List of main conservation measures		
Adapt/manage renewable energy instal	lation, facilities and operation (CC03)	
Reduce impact of multi-purpose hydrological changes (CJ02)		
Implement climate change adaptation measures (CN02)		
Reduce impact of mixed source pollution (CJ01)		
Adapt mowing, grazing and other equivalent agricultural activities (CA05)		
Prevent conversion of (semi-) natural habitats into forests and of (semi-)natural forests into intensive forest plantation (CB01)		
(CB01)		

Adapt/manage exploitation of energy resources (CC02)

Manage water abstraction for resource extraction and energy production (CC13)

Early detection and rapid eradication of invasive alien species of Union concern (CIO1)

8.6 Additional information

#### 9. Future prospects

9.1 Future prospects of parameters

- a) Range
- b) Area
- c) Structure and functions

9.2 Additional information

#### 10. Conclusions

10.1. Range

10.2. Area

10.3. Specific structure and functions

(incl. typical species)

10.4. Future prospects

10.5 Overall assessment of

**Conservation Status** 

10.6 Overall trend in Conservation

Status

10.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

10.8 Additional information

#### 11. Natura 2000 (pSCIs, SCIs, SACs) coverage for Annex I habitat types

11.1 Surface area of the habitat type inside the pSCIs, SCIs and SACs network (in km² in biogeographical/marine region)

11.2 Type of estimate

11.3 Surface area of the habitat type inside the network Method used

11.4 Short-term trend of habitat area in good condition within the network Direction

11.5 Short-term trend of habitat area in good condition within network Method used

11.6 Additional information

a) Minimum

b) Maximum

c) Best single value 5.73

95% confidence interval

Complete survey or a statistically robust estimate

Decreasing (-)

Complete survey or a statistically robust estimate

#### 12. Complementary information

12.1 Justification of % thresholds for trends

12.2 Other relevant information

### **Distribution Map**




Figure 1: UK distribution map for H3130 - Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or of the *Isoëto-Nanojuncetea*. Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available habitat records which are considered to be representative of the distribution within the current reporting period. For further details see the 2019 Article17 UK Approach document.

### Range Map




Figure 2: UK range map for H3130 - Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea uniflorae* and/or of the *Isoëto-Nanojuncetea*. Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by applying a bespoke range mapping tool for Article 17 reporting (produced by JNCC) to the 10km grid square distribution map presented in Figure 1. The alpha value for this habitat was 25km. For further details see the 2019 Article 17 UK Approach document.

### **Explanatory Notes**

Habitat code: 3130 Field label 2.3 Distribution map; Method used 2.3 Short term trend; Shorthern Ireland reports 2008 and 2015 which resurveyed the majority the original H3130 lake locations.  Habitat code: 3130 Region code: ATL Filed label 2.3 Short term trend; Direction 3. The trend of H3130 lake type in Northern Ireland is derived from comparing historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by FNISI Ltd (Environment Science Services) when completing file Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations.  4.5 Short term trend; Method used 3. Distribution of H3130, which was undertaken by FNISI Ltd (Environment Science Services). This completed Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland of H3130, which was undertaken by FNISI Ltd (Environment Science Services). This completed Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  4.9 Long term trend; Method used 3.1 The long-term trend is derived from most comprehensive survey of Northern Ireland of H3130, which was undertaken by FNISI Ltd (Environment Science Services) when completing the Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015. Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015. Which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland reports 2008 and 2015. which resurveyed the majorit		
compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS). The same distribution was recorded by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015 which resurveyed the majority the original H3130 lake locations.  **Mabitat code: 3130 Region code: ATL Field label  **Note**  1		Note
Field label	•	compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS). The same distribution was recorded by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015 which resurveyed the majority the
4.3 Short term trend; Direction  The trend of H3130 lake type in Northern Ireland is derived from comparing historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations.  4.5 Short term trend; Method used  4.5 Short term trend; Method used  The short -term trend is derived from most comprehensive survey of Northern Ireland of H3130, which was undertaken by ENSIS Ltd (Environment Science Services). This completed Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  4.9 Long term trend; Method used  The long-term trend is derived from most comprehensive survey of Northern Ireland of H3130, which was undertaken by ENSIS Ltd (Environment Science Services) to complete Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  5.1 Year or period  The surface area of H3130 lake type in Northern Ireland is derived from comparing historical data compiled by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland is made up of three large lakes - Macnean Upper, Macnean Lower and Lough Melvin, which in total comprise 12.26 km2  5.2 Surface area; Method used  5.5 Short term trend; Period  See 5.1  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes ha	Habitat code: 3130 Region cod	de: ATL
Direction data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations.  4.5 Short term trend; Method used The short -term trend is derived from most comprehensive survey of Northern Ireland of H3130, which was undertaken by ENSIS Ltd (Environment Science Services). This completed Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  4.9 Long term trend; Method used The long-term trend is derived from most comprehensive survey of Northern Ireland of H3130, which was undertaken by ENSIS Ltd (Environment Science Services) to complete Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  5.1 Year or period The surface area of H3130 lake type in Northern Ireland is derived from comparing historical data compiled between 1988 and 1990 by the Northern Ireland reports 2008 and 2015. Which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland is made up of three large lakes - Macnean Upper, Macnean Lower and Lough Melvin, which in total comprise 12.26 km2  5.2 Surface area; Method used See 5.1  5.3 Type of estimate See 5.1  5.4 Surface area; Method used See 5.1  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrien	Field label	Note
of H3130, which was undertaken by ENSIS Ltd (Environment Science Services). This completed Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  4.9 Long term trend; Method used H3130, which was undertaken by ENSIS Ltd (Environment Science Servises) to complete Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  5.1 Year or period The surface area of H3130 lake type in Northern Ireland is derived from comparing historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland is made up of three large lakes - Macnean Upper, Macnean Lower and Lough Melvin, which in total comprise 12.26 km2  5.2 Surface area  5.3 Type of estimate See 5.1  5.4 Surface area; Method used  5.5 Short term trend; Period See 5.1  5.6 Short term trend; Period See 5.1  5.6 Short term trend; Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used	•	data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake
H3130, which was undertaken by ENSIS Ltd (Environment Science Servises) to complete Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015.  5.1 Year or period  The surface area of H3130 lake type in Northern Ireland is derived from comparing historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland is made up of three large lakes - Macnean Upper, Macnean Lower and Lough Melvin, which in total comprise 12.26 km2  5.2 Surface area  5.3 Type of estimate  See 5.1  5.4 Surface area; Method used  See 5.1  5.5 Short term trend; Period  See 5.1  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used		of H3130, which was undertaken by ENSIS Ltd (Environment Science Services). This completed Site Condition Assessments of Standing Water Features in SACs and ASSIs:
historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland is made up of three large lakes - Macnean Upper, Macnean Lower and Lough Melvin, which in total comprise 12.26 km2  5.2 Surface area  see 5.1  5.4 Surface area; Method used  5.5 Short term trend; Period  See 5.1  5.6 Short term trend; Period  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used		H3130, which was undertaken by ENSIS Ltd (Environment Science Servises) to complete Site Condition Assessments of Standing Water Features in SACs and ASSIs: Northern
5.3 Type of estimate  5.4 Surface area; Method used  5.5 Short term trend; Period  5.6 Short term trend;  Direction  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used	5.1 Year or period	historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing the Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed the majority the original H3130 lake locations. Approximately half of the overall resource of the H3130 in Northern Ireland is made up of three large lakes - Macnean Upper, Macnean Lower and Lough Melvin,
5.4 Surface area; Method used  5.5 Short term trend; Period See 5.1  5.6 Short term trend;  Direction  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used	5.2 Surface area	see 5.1
<ul> <li>used</li> <li>5.5 Short term trend; Period See 5.1</li> <li>5.6 Short term trend; Direction Direction Direction See 5.1</li> <li>5.8 Short term trend; Direction Direction See 5.1</li> <li>5.8 Short term trend; Method used</li> </ul>	5.3 Type of estimate	See 5.1
5.6 Short term trend; Direction  Lakes are rarely 'lost' in the conventional sense, although small water bodies may be infilled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used		See 5.1
Direction filled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake types are defined.  5.8 Short term trend; Method used	5.5 Short term trend; Period	See 5.1
used	·	filled or drained. However, many lakes have been severely degraded to the extent that they no longer support characteristic plant or animal communities. As a consequence area assessments show little significant change over time in spite of nutrient enrichment. Degraded sites are not considered lost because of the way in which lake
5.9 Long term trend; Period See 5.1	•	See 5.1
	5.9 Long term trend; Period	See 5.1

5.10 Long term trend; Direction	see 5.6
5.12 Long term trend; Method used	See 5.1
6.1 Condition of habitat	Based on individual waterbody assessments as part of - a) ENSIS Ltd (Environment Science Services) Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland 2008 and 2015, which resurveyed the majority the original H3130 small lake (less 100ha) locations. This identified that only about on-third of the area assessed was in favourable condition; - and b) Water Framework Directive monitoring of Upper Lough Macnean, Lower Lough Macnean and Lough Melvin, which has recorded that all these waterbodies are currently in unfavourable condition.
6.2 Condition of habitat; Method used	see 6.1
6.4 Short term trend of habitat area in good condition; Direction	see 6.1
6.5 Short term trend of habitat area in good condition; Method used	see 6.1
7.1 Characterisation of pressures/ threats	Water pollution - The upland catchments where some of these lakes occur are generally isolated from human occupation but subject to afforestation, resulting in elevated nutrient levels and sedimentation through diffuse pollution resulting from forestry operations. H3130 lakes in mid to low altitude occur in more intensively farmed catchments, which increases the nutrient loading through point and diffuse pollution of surface and groundwater. Sewage - The impact of point source pollution from septic tanks from individual dwellings is recognised as a threat to these low nutrient systems. The increase in aerial nitrogen deposition (APIS) resulting mainly from agricultural activity is now identified as a major threat to all habitats in Northern Ireland. Blanket bog integrity - a proportion of H3130 water bodies occur in the uplands and are often dependent on the condition of the surrounding peatland. Therefore, any adverse impacts on the peatland - such as intensive grazing, mechanical removal of peat, burning and construction of wind farms - which have the potential to induce changes in hydraulic condition in addition to increased sediment loading, may have a negative effect on these upland water bodies. Hence pollution is from multiple sources and reported under J01 Mixed source pollution to surface and ground waters (limnic and terrestrial). Water abstraction - a few of these larger lakes are used as a public water supply. Alien species - the continued spread of invasive aliens is now recognised as a major threat to the freshwater environment. Climate Change - The impact of climate change in the short term is not apparent, however if the predicted changers in temperature and rainfall patterns occur along with extreme weather events there will inevitably impact on the structure and function of all standing freshwater bodies to a greater or lesser degree.

### 8.2 Main purpose of the measures taken

There are two major elements to the current measures being undertaken to restore structure and function. The first is the ongoing commitment to the Water Framework directive (WFD) and carried out through River Basin Management, a key element in implementing the WFD, taking an integrated approach to the protection, improvement and sustainable use of the water environment. It applies to groundwater and to all surface water bodies, including rivers, lakes, transitional (estuarine) and coastal waters out to one nautical mile. In 2009 the first set of River Basin Management Plans (RBMP) as required by the Water Framework Directive were published for each River Basin District within Northern Ireland. The Plan identified where our water environment is in good or excellent conditions and set out objectives for improvement or prevention of deterioration. As required by WFD, the Plans are to be reviewed and updated every 6 years. In 2015 the second set of Plans were published providing an overview of changes and progress that have been made. Northern Ireland has 571 surface and groundwater bodies - 496 surface and 75 groundwater. The 2015 classification results indicate 36.78% are at good or better status. By 2021 with objectives set we aim to increase this to 69.8% with 99.1% at good or better status by 2027. The second is the initiation of the development of Conservation Management Plans for Natura 2000 sites, with a target of 95% completion by December 2020. This project will be undertaken partly through the Rural Development Programme (RDP) - Northern Ireland which will support the development of plans for some Natura 2000 sites, with many of the remaining plans being primarily funded through INTERREG VA. These plans will specifically include Lough Melvin the largest H3130 lake in Northern Ireland, Upper Lough Erne the second largest H3150 lake in Northern Ireland and all six SAC rivers. In addition plans are proposed for all upland peatland SACs which include a significant number of Northern Ireland's small H3130 and H3160 water bodies.

### 9.1 Future prospects of parameters

There are a number of factors which cannot be assessed with any degree of certainty. They include: The effectiveness of the SAC management plans currently being developed, the future impact of aerial nitrogen deposition, climate change and the effectiveness of the EU directives following Brexit, primarily the WFD and HD.

### 11.1 Surface area of the habitat type inside the pSCIs, SCIs and SACs network

The surface area of H3130 feature lakes in the Northern Ireland SAC network is derived from comparing historical data compiled between 1988 and 1990 by the Northern Ireland Lake Survey (NILS) and data compiled by ENSIS Ltd (Environment Science Services) when completing Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed all H3130 feature lakes over 1ha with in the network. The vast majority of the overall SAC area is contributed by Lough Melvin.

#### 11.2 Type of estimate

see 11.1

### 11.3 Surface area of the habitat type inside the network; Method used

see 11.1

# 11.4 Short term trend of habitat area in good condition within the network; Direction

The condition of H3130 feature lakes in the Northern Ireland SAC network is derived from the ENSIS Ltd (Environment Science Services) Site Condition Assessments of Water Features in SACs and ASSIs: Northern Ireland reports 2008 and 2015, which resurveyed all H3130 feature lakes over 1ha within the network, and from WFD monitoring of Lough Melvin.

# 11.5 Short term trend of habitat area in good condition within the network; Method used

see 11.4