European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Supporting documentation for the conservation status assessment for the habitat:

H8120 - Calcareous and calcshist screes of the montane to alpine levels (*Thlaspietea rotundifolii*)

SCOTLAND

IMPORTANT NOTE - PLEASE READ

- The information in this document is a country-level contribution to the UK Report on the conservation status of this habitat, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- The 2019 Article 17 UK Approach document provides details on how this supporting information was used to produce the UK Report.
- The UK Report on the conservation status of this habitat is provided in a separate document.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Explanatory notes (where provided) by the country are included at the end. These provide an audit trail of relevant supporting information.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was only relevant at UK-level (sections 10 Future prospects and 11 Conclusions).
- For technical reasons, the country-level future trends for Range, Area covered by habitat and Structure and functions are only available in a separate spreadsheet that contains all the country-level supporting information.
- The country-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

NATIONAL LEVEL			
1. General information			
1.1 Member State	UK (Scotland information only)		
1.2 Habitat code	8120 - Calcareous and calcshist screes of the montane to alpine levels (Thlaspi		
2. Maps			
2.1 Year or period	1973-2005		

2.1 Year or period	1973-2005
2.3 Distribution map	Yes

2.3 Distribution map Method used Based mainly on extrapolation from a limited amount of data

2.4 Additional maps

BIOGEOGRAPHICAL LEVEL

3. Biogeographical and marine regions

3.1 Biogeographical or marine region where the habitat occurs

Atlantic (ATL)

3.2 Sources of information

References within -

http://jncc.defra.gov.uk/pdf/Article17Consult 20131010/H8120 SCOTLAND.pdf SNH SCM database, extract A2298772, 2017, processed and summarised in A2498677.

Calcareous scree feature type (JNCC, (2009), Common Standards Monitoring Guidance for Upland Habitats, Version July 2009 and previous versions) http://jncc.defra.gov.uk/page-2237

4. Range

- 4.1 Surface area (in km²)
- 4.2 Short-term trend Period
- 4.3 Short-term trend Direction
- 4.4 Short-term trend Magnitude
- 4.5 Short-term trend Method used
- 4.6 Long-term trend Period
- 4.7 Long-term trend Direction
- 4.8 Long-term trend Magnitude
- 4.9 Long-term trend Method used
- 4.10 Favourable reference range

Stable (0)

a) Minimum

b) Maximum

a) Minimum

b) Maximum

- a) Area (km²)
- b) Operator
- c) Unknown No
- d) Method

4.11 Change and reason for change in surface area of range

Improved knowledge/more accurate data

The change is mainly due to: Improved knowledge/more accurate data

4.12 Additional information

NB Range entries and comments are made on the basis of Distribution maps and assumptions as to how these will affect previous range maps, without having seen new range maps. 1) Newly collated vegetation map information (HabMoS) has identified occurrences of this habitat which did not appear in previous Article 17 reporting distribution maps. Some of the new occurrences are outwith the currently-mapped range and would increase the surface area of the range significantly. The location of some of these occurences is credible, adjacent to

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

existing records or where geology and aerial imagery confirm the habitat, but mis-translation of habitat as conforming to this type is also evident. Therefore verification of records would be required before they can be used. NB only a cursory examination of additional occurrences has been possible. 2) For the previously-reported occurrences of the habitat, there is no evidence of any actual change in range in Scotland in the period 2006-2017. Within this period, persistence of the habitat has been confirmed in all the upland designated sites where it is a notified feature that have been checked (SCM database, extract A2298772).

the period. Within this period, no losses of extent have been recorded on sites where it is a notified feature (SCM database, extract A2298772). Vegetation overgrowth on scree can potentially lead to some losses of extent; increases in

5. Area covered by habitat

5.1 Year or period	2007-007-		
5.2 Surface area (in km²)	a) Minimum 1	b) Maximum 2	c) Best single 1.5 value
5.3 Type of estimate	Best estimate		
5.4 Surface area Method used	Based mainly on e	xpert opinion with very limite	ed data
5.5 Short-term trend Period	2007-2017		
5.6 Short-term trend Direction	Stable (0)		
5.7 Short-term trend Magnitude	a) Minimum	b) Maximum	c) Confidence interval
5.8 Short-term trend Method used	Based mainly on e	xtrapolation from a limited a	mount of data
5.9 Long-term trend Period			
5.10 Long-term trend Direction			
5.11 Long-term trend Magnitude	a) Minimum	b) Maximum	c) Confidence interval
5.12 Long-term trend Method used			
5.13 Favourable reference area	a) Area (km²)		
	b) Operator		
	c) Unknown N	0	
	d) Method		
5.14 Change and reason for change	Improved knowled	dge/more accurate data	
in surface area of range	The change is main	nly due to: Improved know	wledge/more accurate data
5.15 Additional information	Conclusions are ba	ased on absence of evidence	of change in extent in Scotland in

6. Structure and functions			
6.1 Condition of habitat	a) Area in good condition (km²)	Minimum 0.90909	Maximum 1.81818
	b) Area in not-good condition (km²)	Minimum 0.0909	Maximum 0.18181
	c) Area where condition is not known (km²)	Minimum 0	Maximum 0
6.2 Condition of habitat Method used	Complete survey or a statist	cically robust estimate	

extent through new scree shoots have been observed.

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

6.3 Short-term trend of habitat area in good condition Period

6.4 Short-term trend of habitat area in good condition Direction

6.5 Short-term trend of habitat area in good condition Method used

6.6 Typical species

6.7 Typical species Method used

6.8 Additional information

2003-2014

Stable (0)

Complete survey or a statistically robust estimate

Has the list of typical species changed in comparison to the previous No reporting period?

Site Condition Monitoring provides a means of assessing the structure and function of H8120 in Scotland. Assessment is based on the results of assessments carried out between 2003 and 2014. Results are recorded on the SNH SCM database, from which data was extracted to A2298772 on 23/05/2017. Within this period, the proportion of H8120 on SACs considered to be in Favourable condition remained constant at 90% between 2012 (based on assessments carried out between 2003 and 2010) and 2014. No H8120 is reported as recovering or declining, as in 2012. One SSSI feature not overlapping SAC is considered to be Favourable and none Unfavourable, but extent data is not available. Overall, no habitat was assessed as declining in condition (Unfavourable declining or Favourable declining) or recovering (Favourable recovered, Unfavourable recovering, Unfavourable recovering due to management), the same as in 2012. The proportion in Favourable condition has remained high and stable, and no habitat is assessed as recovering or declining, therefore overall condition is judged to be stable.

7. Main pressures and threats

7.1 Characterisation of pressures/threats

Pressure	Ranking
Management of fishing stocks and game (G08)	M
Intensive grazing or overgrazing by livestock (A09)	M
Threat	Ranking
Management of fishing stocks and game (G08)	M
Intensive grazing or overgrazing by livestock (A09)	M

7.2 Sources of information

7.3 Additional information

Deer grazing and trampling Trampling and grazing by sheep

8. Conservation measures

8.1 Status of measures	a) Are measures needed?	Yes	
	b) Indicate the status of measures	Measures identified and taken	
8.2 Main purpose of the measures taken	Restore the habitat of the species (re	elated to 'Habitat for the species')	
8.3 Location of the measures taken	Only inside Natura 2000		
8.4 Response to the measures	Medium-term results (within the next two reporting periods, 2019-2030)		
8.5 List of main conservation measures			

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

Adapt mowing, grazing and other equivalent agricultural activities (CA05)

Management of hunting, recreational fishing and recreational or commercial harvesting or collection of plants (CG02)

8.6 Additional information

Conservation measures are generally implemented through designation of protected areas, voluntary and statutory procedures (Deer Act), agrienvironment schemes (SRDP). While some results are achievable in the short term, some attributes will recover only over longer timescales.

9. Future prospects

- 9.1 Future prospects of parameters
- a) Range
- b) Area
- 9.2 Additional information
- c) Structure and functions

Range is considered likely to remain stable. Area is considered likely to remain stable. The very high proportion of features Favourable for Structure and function should continue to be so. Condition is good and stable.

10. Conclusions

- 10.1. Range
- 10.2. Area
- 10.3. Specific structure and functions
- (incl. typical species)
- 10.4. Future prospects
- 10.5 Overall assessment of
- **Conservation Status**
- 10.6 Overall trend in Conservation

Status

10.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

10.8 Additional information

11. Natura 2000 (pSCIs, SCIs, SACs) coverage for Annex I habitat types

- 11.1 Surface area of the habitat type inside the pSCIs, SCIs and SACs network (in km² in biogeographical/marine region)
- 11.2 Type of estimate
- 11.3 Surface area of the habitat type inside the network Method used
- a) Minimum
- b) Maximum
- c) Best single value 3.33

Best estimate

Based mainly on extrapolation from a limited amount of data

Report on the main results of the surveillance under Article 17 for Annex I habitat types (Annex D)

11.4 Short-term trend of habitat area in good condition within the network Direction

11.5 Short-term trend of habitat area in good condition within network Method used

11.6 Additional information

Stable (0)

Complete survey or a statistically robust estimate

Site Condition Monitoring provides a means of assessing the structure and function of H8210 on SACs in Scotland. Assessment is based on the results of assessments carried out between 2003 and 2014. Results are recorded on the SNH SCM database, from which data was extracted to A2298772 on 23/05/2017. Within this period, the proportion of H8120 on SACs considered to be in Favourable condition remained constant at 90% between 2012 (based on assessments carried out between 2003 and 2010) and 2014. No H8120 is reported as recovering or declining, as in 2012. Overall, no habitat was assessed as declining in condition (Unfavourable declining or Favourable declining) or recovering (Favourable recovered, Unfavourable recovering, Unfavourable recovering due to management), the same as in 2012. The proportion in Favourable condition has remained high and stable, and no habitat is assessed as recovering or declining, therefore overall condition is judged to be stable.

12. Complementary information

12.1 Justification of % thresholds for trends

12.2 Other relevant information

Distribution Map




Figure 1: UK distribution map for H8120 - Calcareous and calcshist screes of the montane to alpine levels (*Thlaspietea rotundifolii*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available habitat records which are considered to be representative of the distribution within the current reporting period. For further details see the 2019 Article17 UK Approach document.

Range Map




Figure 2: UK range map for H8120 - Calcareous and calcshist screes of the montane to alpine levels (*Thlaspietea rotundifolii*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by applying a bespoke range mapping tool for Article 17 reporting (produced by JNCC) to the 10km grid square distribution map presented in Figure 1. The alpha value for this habitat was 25km. For further details see the 2019 Article 17 UK Approach document.