# European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

# Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Conservation status assessment for the species:

S1096 - Brook lamprey (Lampetra planeri)

**UNITED KINGDOM** 

#### **IMPORTANT NOTE - PLEASE READ**

- The information in this document represents the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- It is based on supporting information provided by the geographically-relevant Statutory Nature Conservation Bodies, which is documented separately.
- The 2019 Article 17 UK Approach document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Maps showing the distribution and range of the species are included (where available).
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the UK assessments. Further underpinning explanatory notes are available in the related country-level reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species).
- The UK-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

NATIONAL LEVEL		
1. General information		
1.1 Member State	UK	
1.2 Species code	1096	
1.3 Species scientific name	Lampetra planeri	
1.4 Alternative species scientific name		
1.5 Common name (in national language)	Brook lamprey	

### 2. Maps

2.1 Sensitive species	No
2.2 Year or period	1990-2018
2.3 Distribution map	Yes
2.4 Distribution map Method used	Based mainly on extrapolation from a limited amount of data
2.5 Additional maps	No

#### 3. Information related to Annex V Species (Art. 14)

5. Illiorination related to 7	Alliex v Species (Alt. 14)	
3.1 Is the species taken in the wild/exploited?	No	
3.2 Which of the measures in Art. 14 have been taken?	a) regulations regarding access to property	No
	b) temporary or local prohibition of the taking of specimens in the wild and exploitation	No
	c) regulation of the periods and/or methods of taking specimens	No
	d) application of hunting and fishing rules which take account of the conservation of such populations	No
	e) establishment of a system of licences for taking specimens or of quotas	No
	f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens	No
	g) breeding in captivity of animal species as well as artificial propagation of plant species	No

h) other measures

No

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

#### a) Unit

b) Statistics/ quantity taken	Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period					
	Season/ year 1	Season/ year 2	Season/ year 3	Season/ year 4	Season/ year 5	Season/ year 6
Min. (raw, ie. not rounded)						
Max. (raw, ie. not rounded)						
Unknown	No	No	No	No	No	No

- 3.4. Hunting bag or quantity taken in the wild Method used
- 3.5. Additional information

#### **BIOGEOGRAPHICAL LEVEL**

#### 4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

#### 4.2 Sources of information

Atlantic (ATL)

England

Addy, S., Cooksley, S., Dodd, N., Waylen, K., Stockan, J., Byg, A. & Holstead, K. 2016. River restoration and biodiversity: Nature based solutions for restoring rivers in the UK and Republic of Ireland. CREW ref. CRW2014/10

APEM 2005. Review of information on lamprey populations in the Humber basin. Report to the Environment Agency.

APEM 2014. A habitat survey for spawning ground and nursery areas for Annex II fish species within the Severn Estuary river catchment. APEM Scientific Report 413588. Natural England, January 2015, 45pp.

Bracken, F.S.A. & Lucas, M.C. 2013. Potential impacts of small-scale hydroelectric power generation on downstream moving lamprey. River research and applications. 29. 1073-1081.

Bracken, F.S.A., Hoelzel, A.R., Hume, J.B. & Lucas, M.C. 2015. Contrasting population genetic structure among freshwater-resident and anadromous lampreys: the role of demographic history, differential dispersal and anthropogenic barriers to movement.

Common Standards Monitoring Guidance for Freshwater Fauna 2015 Common Standards Monitoring Guidance for Rivers 2014

Davies, C., Shelley, J., Harding, P., McLean, I., Gardiner, Ross & Peirson, G. 2004. Freshwater Fishes in Britain. The species and their distribution. Harley Books. Environment Agency, 2013. Review of fish population data in the Humber Estuary.

Environment Agency, 2012. Summary of outcomes of the Review of Consents on water-related SACs. Excel spreadsheet.

Foulds, L.W. 2013. Anthropogenic factors affecting European river lamprey

Lampetra fluviatilis in the Humber River basin, north-east England. PhD. Thesis, Durham University.

Foulds, L.W. & Lucas, M.C. 2013. Extreme inefficiency of two conventional technical fishways used by European river lamprey (Lampetra fluviatilis). Ecological Engineering, 58, 423-433.

Foulds, W.L. & Lucas, M.C. 2014. Paradoxical exploitation of protected fishes as bait for anglers: Evaluating the lamprey bait market in Europe and developing sustainable and ethical solutions. PLOS ONE, Vol. 9, Issue 6, e99617.

Findlay, J.D.S. 2013. Impacts of signal crayfish on stream fishes. Durham theses, Durham University.

Findlay, J.D.S., Riley, W.D. & M.C. Lucas. 2014. Signal crayfish (Pacifastacus leniusculus) predation upon Atlantic salmon (Salmo salar) eggs. Aquatic Conservation: Marine and Freshwater Ecosystems. 25. 250-258.

Guan, R-Z. & Wiles, P.R. 1996. Ecological Impact of Introduced Crayfish on Benthic Fishes in a British Lowland River. Conservation Biology, 11

Greaves, R.K., Bubb, D.H. & Lucas, M.C. 2007. Adult river lamprey occurrence and migration in the River Trent in relation to barriers and environmental conditions 2006-2007. Final report. University of Durham.

Harvey, J. & Cowx I. 2003. Monitoring the River, Brook and Sea Lamprey, Lampetra fluviatilis, L. planeri and Petromyzon marinus. Conserving Natura 2000 Rivers Monitoring Series No. 5, English Nature.

Hatton-Ellis, T. 2018. Procedure for Estimating Population (including Favourable Reference Population) using 1km Square Resolution Records Data. Interagency freshwater group. (Unpublished).

Holdich, D.M., James, J., Jackson, C. & Peay, S. 2014. The North American signal crayfish, with particular reference to its success as an invasive species in Great Britain. Ethology, Ecology & Evolution, 26, 232-262.

Jacklin, T.E. 2006. The status of lamprey species in the River Trent with particular regard to the Humber Estuary SAC. Environment Agency.

Langford, T.E., Worthington, T., Shaw, P., Kemp, P., Woolgar, C., Fergusson, A., Harding, P & Ottewell, D. 2012. The unnatural history of the River Trent: 50 years of ecological recovery. River Conservation and Management. Boon, J.P. & Raven, P.J. (Eds.). John Wiley & Sons, Ltd.

Lucas, M. & Bracken, F. 2010. Potential impacts of hydroelectric power generation on downstream-moving lampreys at Howsham, Yorkshire Derwent. Durham University. (Unpublished).

Mainstone, C.P., Dils, R.M. and Withers, P.J.A. 2008. Controlling sediment and phosphorus transfer to receiving waters - A strategic management perspective for England and Wales. Journal of Hydrology, 350, 131-143.

Mainstone, C.P. and Holmes, N.T. 2010. Embedding a strategic approach to river restoration in operational management processes - experiences in England. Aquatic Conservation: Marine and Freshwater Ecosystems. Published online in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/aqc.1095 Mainstone, C., Hall, R. & Diak, I. 2016. A narrative for conserving freshwater and wetland habitats in England. Natural England Research Reports, Number 064. Mainstone C.P. 2008. The role of specially designated wildlife sites in freshwater conservation - an English perspective. Freshwater Reviews, 1, 89-98.

Mainstone, C. & Burn, A. 2011. Relationships between ecological objectives and associated decision-making under the Habitats and Water Framework Directives. Discussion paper, Natural England.

Mainstone, C.P. & Wheeldon, J. 2016. The physical restoration of English rivers with special designations for wildlife: from concepts to strategic planning and implementation. Freshwater Reviews. 8. Pg. 1 - 25.

Mainstone, C.P. 2016. Developing a coherent narrative for conserving freshwater and wetland habitats: experiences in the UK. WIRES Water, published Online: Nov 07 2016. DOI: 10.1002/wat2.1189.

Mainstone, C.P. 2018. Article 17 Habitats Pro-forma England H3260 for UK aggregation. Natural England

Mainstone, C.P. 2018. Analysis of Water Framework Directive data for use in Habitats Directive Article 17 reporting on Annex I river habitat (H3260) in England. Supplementary paper for the submission package to Europe, Natural England.

Maitland, P.S. 1980. Review of the ecology of lampreys in northern Europe. Canadian Journal of Fisheries and Aquatic Sciences. 37, 1944-1952.

Maitland, P.S. 2003. Ecology of the River, Brook and Sea Lamprey. Conserving Natura 2000 Rivers, Ecology Series No. 5. English Nature, Peterborough Maria, I., Bernardo, J.M. & Fernandes, S. 2007. Predation of invasive crayfish on aquatic vertebrates: the effect of Procambarus clarkii on fish assemblages in Mediterranean temporary streams. Biological Invaders in Inland Waters. Gherardi, F. (ed.)

Natural England. 2015. River restoration theme plan. Output from the EU Life project 'Improvement Programme for England's Natura 2000 Sites' (IPENS). Natural England Report number IPENSTP023.

Nunn, A.D., Harvey, J.P., Noble, R.A.A. & Cowx, I.G 2008. Condition assessment of lamprey populations in the Yorkshire Ouse catchment, north-east England, and the potential influence of physical migration barriers. Aquatic Conservation: Marine and Freshwater Ecosystems 18: 175-189.

Nunn, A.D., Harvey, J.P., Noble, R.A.A. & Cowx, I.G. 2015. A desk-based study to consolidate existing data and literature to help inform the conservation status of Annex II features (migratory fish species) within the Severn Estuary river catchment. Natural England, IPENS programme LIFE11NAT/UK/000384IPENS. Peay, S., Guthrie, N., Spees, J., Nilsson, E. & Bradley, P. 2009. The impact of signal crayfish (Pacifastacus leniusculus) on the recruitment of salmonid fish in a headwater stream in Yorkshire, England. Knowledge and Management of Aquatic Ecosystems.

Reynolds, J.D. 2011. A review of ecological interactions between crayfish and fish, indigenous and introduced. Knowledge and Management of Aquatic Ecosystems. 401, 10

Russon, I.G., Kemp, P.S. & Lucas, M.C. 2011. Guaging weirs impede the upstream migration of adult river lamprey Lampetra fluviatilis. Fisheries Management and Ecology. 18. 201-2010.

Tweed Estuary SAC / River Tweed SAC. Assessment of lamprey spawning habitat in the Till catchment in England. 2015. Report produced by The Tweed Foundation for Natural England.

Wheeldon, J. 2018. Progress report on the English river SSSI/SAC physical restoration programme. Paper to the river SSSI restoration project steering group.

Scotland

Ecological Research Associates. 2017. National Lamprey Survey of Scotland. Scottish Natural Heritage Commissioned Report No. 464.

Forth Fisheries Foundation. 2004. River and brook lamprey monitoring of the Endrick Water cSAC/SSSI. Scottish Natural Heritage Commissioned Report No. 057.

Watt J, Bull C, Ravenscroft, NOM, Seed M. 2011. Lamprey Survey of the Endrick Water SSSI/SAC 2008. Scottish Natural Heritage Commissioned Report No. 320. Bull C, Perfect C, Watt J. 2016. Site condition monitoring of lamprey in the

Endrick Water SSSI and SAC 2012. Scottish Natural Heritage Commissioned Report No. 911. https://www.nature.scot/snh-commissioned-report-911-site-condition-monitoring-lamprey-endrick-water-sssi-and-sac-2012

Hume JB. 2011 Adult lamprey survey of the Endrick Water SSSI and SAC 2009-2010. Scottish Natural Heritage Commissioned Report No. 480.

Watt, J., Brown, L. & Bull, C. 2011. Lamprey Site Condition Monitoring of the River Spey SSSI and SAC. Unpublished Scottish Natural Heritage commissioned report.

APEM. 2004. Distribution of sea, brook and river lampreys on the River Tay. Scottish Natural Heritage Commissioned Report No. 032.

Watt J, Ravenscroft NOM, Seed M. 2008. Site Condition Monitoring of lamprey in the River Tay SAC. Scottish Natural Heritage Commissioned Report No. 292.

Watt J, Bull C. Site Condition Monitoring of lamprey in the River Tay SAC 2012-2013. Unpublished Scottish Natural Heritage commissioned report.

Maitland PS. 2000. Distribution of lampreys in the River Teith. Unpublished Scottish Natural Heritage commissioned report.

Bull C. 2003. Electro-fishing survey in selected tributaries of the River Teith candidate Special Area of Conservation. Unpublished Scottish Natural Heritage commissioned report.

Bull C, Watt J. Site condition monitoring of lamprey in the River Teith SAC 2011. Unpublished Scottish Natural Heritage commissioned report.

Bull C. Site condition monitoring of lamprey in the River Teith SAC 2015-2016. Unpublished Scottish Natural Heritage commissioned report.

Bull C, Shelton J, Coulson M. 2018. The collection of larval sea lamprey (Petromyzon marinus) using drift netting, suction pumping, and airlift sampling: a trial to inform the development of methods suitable for Site Condition Monitoring. Scottish Natural Heritage Research Report No. 1040.

The Tweed Foundation. 2004. Assessment of lamprey distribution and abundance in the River Tweed cSAC/SSSI. Unpublished Scottish Natural Heritage commissioned report.

IAFG. 2017. UK Article 17 reporting procedure for estimating population using 1 km square resolution records data. Inter-agency Freshwater Group, UK. Watt J, Bull C, Ravenscroft NOM. Site Condition Monitoring of lamprey in the River Tweed SSSI and SAC 2011. Unpublished Scottish Natural Heritage commissioned report.

Records from the Atlas of Freshwater Fishes accessed through NBN Atlas website (https://registry.nbnatlas.org/public/show/dr741).

Records from the Freshwater Fish Recording Scheme accessed through the NBN Atlas website (https://registry.nbnatlas.org/public/show/dr1851).

North East Scotland Fish Records 1800-2010 accessed through the NBN Atlas website (https://registry.nbnatlas.org/public/show/dr1230).

NBN Atlas website at http://www.nbnatlas.org accessed 29 June 2018. Wales

Angling Times. 2017. The best dead baits to use when angling for pike. Dead baiting is the number one pike tactic in the UK.

https://www.anglingtimes.co.uk/advice/bait/articles/the-best-deadbaits-to-use-when-pike-fishing [Accessed 10 April 2018]

Baxter E, McKenzie S, Jones C, Jones D. & Metcalfe P. 2017. Condition assessment using 2016 River Habitat Survey data and Common Standards Monitoring guidance for the Afon Teifi and Afon Eden - Cors Goch Trawsfynydd SACs. NRW Evidence Report No: 192, 92 pp. NRW, Bangor.

Blank M, Jurss K, Bastrop E. 2008. A mitochondrial multigene approach contributing to the systematics of the brook and river lampreys and the

phylogenetic position of Eudontomyzon mariae. Canadian Journal of Fisheries and Aquatic Sciences. 65(12): 2780-279.

Bracken FSA, Hoelzel AR, Hume JB, Lucas MC. 2015. Contrasting population genetic structure among freshwater-resident and anadromous lampreys: the role of demographic history, differential dispersal and anthropogenic barriers to movement. Molecular Ecology, 24: 1188-1204.

Campbell D, Williams E, APEM Aquatic Scientists. 2006. Lamprey Survey on the River Dee and Tributaries: Final Report - March 2006. Environment Agency Wales. Dawson H, Quintella B, Almeida P, Treble A, Jolley J. 2015. The Ecology of Larval and Metamorphosing Lampreys. In: Docker M. (eds) Lampreys: Biology, Conservation and Control. Fish & Fisheries Series, vol 37. Springer, Dordrecht. Espanhol R, Almeida P, Alves JM. 2007. Evolutionary history of lamprey paired species Lampetra fluviatilis (L.) and Lampetra planeri (Bloch) as inferred from mitochondrial DNA variation. Molecular Ecology,16. 1909-1924.

Gardiner R. 2003. Identifying Lamprey. Conserving Natura 2000 Rivers Techniques Series No. 4. Peterborough, English Nature.

Garrett H, Thomas Rh. 2012. Afon Tywi Population Attribute Condition Assessment for Brook, River and Sea Lamprey 2011. CCW Staff Science Report No. 11/8/5. Bangor, Countryside Council for Wales.

Garrett H, Hatton-Ellis TW, Thomas Rh. 2013a. Afonydd Cleddau Population Attribute Condition Assessment for Brook, River and Sea Lamprey 2012. CCW Staff Science Report No. 13/8/1. Bangor: Countryside Council for Wales. Garrett H, Thomas Rh, Hatton-Ellis TW. 2013b. River Usk Population Attribute Condition Assessment for Brook, River and Sea Lamprey 2007-12. CCW Staff Science Report No. 11/8/6. Bangor: Countryside Council for Wales.

Garrett HM. 2016. River Usk SAC habitat structure condition assessment using 2013 - 2015 RHS data & Common Standards Monitoring guidance. NRW Evidence Report No 142, 28, Dolgellau: NRW.

Garrett HM, Thomas Rh. 2016. River Wye SAC habitat structure condition assessment using 2013 - 2015 RHS data & Common Standards Monitoring guidance. NRW Evidence Report No 141, 28pp. Dolgellau: NRW.

Garrett HM. 2015. River Dee & Bala lake SAC population condition attribute condition assessment for brook, river and sea lamprey population 2014. NRW Evidence Report No: 40 31pp, Dolgellau: NRW.

Garrett HM. 2017. SAC monitoring summary note. River Wye SAC lamprey species population condition assessment. Reporting cycle 2013 - 2018. Bangor: NRW.

Garrett, HM. 2016. Afon Teifi SAC population attribute condition assessment for brook, river and sea lamprey population 2014. NRW Evidence Report No. 106. 28 pp. Bangor: NRW.

Hardisty M. 2006. Lampreys. Life without jaws. Pub. Forrest text. Interagency Freshwater Group (IAFG). 2017. UK Article 17 reporting. Procedure for estimating population (inc favourable Reference Population) using 1km square resolution records data. December 2017. Interagency Freshwater Group. Joint Nature Conservation Committee (JNCC). 2015. Common Standards Monitoring Guidance for Freshwater Fauna. Peterborough, Joint Nature Conservation Committee.

JNCC. 2007. Second Report by the UK under Article 17 on the implementation of the Habitats Directive from January 2001 to December 2006. Peterborough: JNCC. Available from: www.jncc.gov.uk/article17.

Living with Environment Change (LWEC). 2011-2015. Climate change impact report cards. Available online:

http://www.nerc.ac.uk/research/partnerships/lwec/products/report-cards/.

Accessed January 2018.

Maitland PS. 2003. Ecology of the River, Brook and Sea Lamprey. Conserving Natura 2000 Rivers Ecology Series No. 5. English Nature, Peterborough. Mateus CS, Almeida PR, Mesquita N, Quintella BR, Alves MJ. 2016. European lampreys: new insights on postglacial colonization, gene flow and speciation. PLoS ONE 11(2): e0148107.

Mateus CS, Stange M, Berner D, Roesti M, Quintella BR, Alves MJ, Almeida PR, Salzburger W. 2013. Strong genome-wide divergence between sympatric European river and brook lampreys. Curr Biol. 23(15): R649-50.

Natural Resources Wales. 2013. Supporting documentation for the Third Report by the United Kingdom under Article 17 on the implementation of the Directive from January 2007 to December 2012 Conservation status assessment for Species: S1096 - Brook lamprey (Lampetra planeri).

[http://jncc.defra.gov.uk/pdf/Article17Consult\_20131010/S1096\_WALES.pdf] Natural Resources Wales. 2015. Water Watch Wales maps gallery. Cycle 2 waterbodies and rivers.

https://nrw.maps.arcgis.com/apps/webappviewer/index.html?id=2176397a06d6 4731af8b21fd69a143f6

Natural Resources Wales. 2017. National Fish Populations database held on BIOSYS. Accessed December 2017.

Schreiber, A & Engelhorn S. 1998. Population genetics of a cyclostome species pair, river lamprey (Lampetra fluviatilis L.) and brook lamprey (Lampetra planeri Bloch). Journal of Zoological Systematics and Evolutionary Research, 36, Issue1-2, 85-99.

Teague N, Webb H, Allen V, Cesar CP, Thomas Rh, Hatton-Ellis TW. 2012. Lamprey monitoring on the River Dee special area of conservation SAC 2011. CCW Science Report No. 975. Bangor: Countryside Council for Wales (CCW). Wilson L, McCall R, Astbury S, Bhogal A, Walmsley C. 2013. Climate Vulnerability Assessment of Designated Sites in Wales. CCW Contract Science Report No. 1017. CCW. Bangor.

N.Ireland

Goodwin, C (2003). Ecology of three lamprey species in Northern Ireland. Goodwin, C.E., Dick, J.T.A., Rogowski, D.L., Elwood, R.W. (2008). Lamprey ammocoete habitat association at regional, catchment and microhabitat scales in Northern Ireland. Ecology of Freshwater Fish. Blackwell Munksgaard: 17: 542-553.

Loughs Agency (2010). Lamprey Baseline Surveys: River Finn and River Deele Co Donegal. Loughs Agency of the Foyle Carlingford and Irish Lights Commission. Report Ref: LA/Lamprey/04&09/11. https://www.loughs-agency.org/wp-content/uploads/2015/05/lamprey-baseline-surveys-finn-and-deele-2010.pdf. Maitland PS (2003). Ecology of the River, Brook and Sea Lamprey. Conserving Natura 2000 Rivers Ecology Series No. 5. English Nature, Peterborough. Niven, A & McCauley (2013a). Lamprey baseline survey No. 2: River Faughan and Tributaries SAC. Loughs Agency, L'Derry.

Niven, A & McCauley (2013b). Lamprey baseline survey No. 3: River Foyle and Tributaries SAC. Loughs Agency, L'Derry.

Niven, A & McCauley (2013c). Lamprey baseline survey No. 4: River Roe and Tributaries SAC. Loughs Agency, L'Derry.

Kurz, I and Costello, M.J. 1999 An outline of the biology, distribution and conservation of Lampreys in Ireland. Irish Wildlife Manuals No 5. Ducjas, Dublin.

#### 5. Range

5.1 Surface area (km²) 169139.04 5.2 Short-term trend Period 2007-2018 5.3 Short-term trend Direction Stable (0) 5.4 Short-term trend Magnitude b) Maximum a) Minimum 5.5 Short-term trend Method used Based mainly on extrapolation from a limited amount of data 5.6 Long-term trend Period 5.7 Long-term trend Direction 5.8 Long-term trend Magnitude a) Minimum b) Maximum 5.9 Long-term trend Method used 5.10 Favourable reference range a) Area (km²) 168863 b) Operator c) Unknown d) Method The FRR is the same as in 2013. The value is considered to be large enough to support a viable population and no lower than the range estimate from when the Habitats Directive came into force in the UK. For further details see the 2019 Article 17 UK Approach document. 5.11 Change and reason for change Improved knowledge/more accurate data in surface area of range Use of different method The change is mainly due to: Use of different method 5.12 Additional information It is not possible to distinguish between Brook and River lamprey juvenilles (amnocetes), so many records are at the Lampetra genus level. Therefore, there is inherent uncertainty in the Brook lamprey distribution, in England the distribution is thought to be under-represented. 6. Population 6.1 Year or period 1990-2018 6.2 Population size (in reporting unit) a) Unit number of map 1x1 km grid cells (grids1x1) b) Minimum c) Maximum d) Best single value 1865 6.3 Type of estimate Minimum 6.4 Additional population size (using a) Unit population unit other than reporting b) Minimum unit) c) Maximum d) Best single value 6.5 Type of estimate 6.6 Population size Method used Based mainly on extrapolation from a limited amount of data 6.7 Short-term trend Period 2006-2018 6.8 Short-term trend Direction Unknown (x)

ii, iv aliu v species (Alii	iex bj		
6.9 Short-term trend Magnitude	<ul><li>a) Minimum</li><li>b) Maximum</li><li>c) Confidence interval</li></ul>		
6.10 Short-term trend Method used	Insufficient or no data available		
6.11 Long-term trend Period			
6.12 Long-term trend Direction			
6.13 Long-term trend Magnitude	<ul><li>a) Minimum</li><li>b) Maximum</li><li>c) Confidence interval</li></ul>		
6.14 Long-term trend Method used			
6.15 Favourable reference population (using the unit in 6.2 or 6.4)	<ul><li>a) Population size</li><li>b) Operator</li><li>c) Unknown</li></ul>	x	
	d) Method	The FRP for this species is insufficient information to further information see the Approach document.	set an FRP value. For
6.16 Change and reason for change in population size	Improved knowledge/more accurate data Use of different method		
	The change is mainly d	lue to: Use of different me	ethod
6.17 Additional information	population size and tre distinguish between ri stage. Although ammo	enough information for this sends, and, therefore to set an ver lamprey and brook lamp accete density measures are e identified to species.	n FRP. It is not possible to rey at the ammocoetes life
7. Habitat for the species			
7.1 Sufficiency of area and quality of occupied habitat	a) Are area and quality sufficient (for long-term	•	Unknown
	b) Is there a sufficientl habitat of suitable qua survival)?	y large area of unoccupied lity (for long-term	Unknown
7.2 Sufficiency of area and quality of occupied habitat Method used	Based mainly on exper	t opinion with very limited o	lata
7.3 Short-term trend Period	2001-2018		
7.4 Short-term trend Direction	Unknown (x)		
7.5 Short-term trend Method used	Insufficient or no data	available	
7.6 Long-term trend Period			
7.7 Long-term trend Direction			
7.8 Long-term trend Method used			

7.9 Additional information

10

Brook lamprey have complex habitat requirements as their eggs are laid in well oxygenated gravels, the ammocoetes require soft sediments and upstream

migratory passage, with relatively low flow velocities and turbulence, must be maintained for adults. These factors, combined with the widespread distribution of brook lamprey, make a detailed assessment of habitat quality trends impossible at the present time

#### 8. Main pressures and threats

#### 8.1 Characterisation of pressures/threats

or pressures, an each	
Pressure	Ranking
Agricultural activities generating point source pollution to surface or ground waters (A25)	M
Agricultural activities generating diffuse pollution to surface or ground waters (A26)	M
Hydropower (dams, weirs, run-off-the-river), including infrastructure (D02)	M
Mixed source pollution to surface and ground waters (limnic and terrestrial) (J01)	M
Modification of hydrological flow (K04)	M
Physical alteration of water bodies (K05)	M
Droughts and decreases in precipitation due to climate change (NO2)	M
Change of habitat location, size, and / or quality due to climate change (N05)	M
Threat	Ranking
Agricultural activities generating point source pollution to surface or ground waters (A25)	M
Agricultural activities generating diffuse pollution to surface or ground waters (A26)	M
Hydropower (dams, weirs, run-off-the-river), including infrastructure (D02)	M
Invasive alien species of Union concern (I01)	M
Mixed source pollution to surface and ground waters (limnic and terrestrial) (J01) $$	M
Modification of hydrological flow (K04)	M
Physical alteration of water bodies (K05)	M
Droughts and decreases in precipitation due to climate change (N02)	M
Change of habitat location, size, and / or quality due to climate change (N05)	M
Other climate related changes in abiotic conditions (N09)	M

8.2 Sources of information

8.3 Additional information

#### 9. Conservation measures

9.1 Status of measures a) Are measures needed? Yes

> b) Indicate the status of measures Measures identified and taken

9.2 Main purpose of the measures Maintain the current range, population and/or habitat for the species

9.3 Location of the measures taken Both inside and outside Natura 2000

9.4 Response to the measures Medium-term results (within the next two reporting periods, 2019-2030)

9.5 List of main conservation measures

Reduce impact of hydropower operation and infrastructure (CC04)

Reduce/eliminate point source pollution to surface or ground waters from industrial, commercial, residential and recreational areas and activities (CF04)

Manage water abstraction for public supply and for industrial and commercial use (CF11)

Management of professional/commercial fishing (including shellfish and seaweed harvesting) (CG01)

Management of hunting, recreational fishing and recreational or commercial harvesting or collection of plants (CG02)

Reduce impact of mixed source pollution (CJ01)

Reduce impact of multi-purpose hydrological changes (CJ02)

Restore habitats impacted by multi-purpose hydrological changes (CJ03)

Adopt climate change mitigation measures (CN01)

Implement climate change adaptation measures (CN02)

9.6 Additional information

#### 10. Future prospects

10.1 Future prospects of parameters a) Range Good

> Unknown b) Population Unknown

c) Habitat of the species

#### 10.2 Additional information 11. Conclusions

11.1. Range Favourable (FV)

11.2. Population Unknown (XX)

11.3. Habitat for the species Unknown (XX)

Unknown (XX) 11.4. Future prospects

11.5 Overall assessment of **Conservation Status** 

11.6 Overall trend in Conservation Status

Unknown (XX)

11.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No information on nature of change

The change is mainly due to:

b) Overall trend in conservation status

No information on nature of change

The change is mainly due to:

11.8 Additional information

Conclusion on Range reached because: (i) the short-term trend direction in Range surface area is stable; and (ii) the current Range surface area is not less than the Favourable Reference Range.

Conclusion on Population reached because: (i) the short-term trend direction in Population size is unknown; and (ii) the Favourable Reference Population is unknown.

Conclusion on Habitat for the species reached because: (i) the area of occupied and unoccupied habitat is unknown and (ii) the habitat quality is unknown for the long-term survival of the species; and (iii) the short-term trend in area of habitat is unknown.

Conclusion on Future prospects reached because: (i) the Future prospects for Range are good; (ii) the Future prospects for Population are unknown; and (iii) the Future prospects for Habitat for the species are unknown.

Overall assessment of Conservation Status is Unknown because three of the conclusions are Unknown.

Overall trend in Conservation Status is based on the combination of the short-term trends for Range - stable, Population - unknown, and Habitat for the species - unknown.

The Overall assessment of Conservation Status has changed between 2013 and 2019 because the conclusion for Habitat for the species has changed from Favourable to Unknown and the conclusion for Future Prospects has changed from Favourable to Unknown.

The Overall trend in Conservation Status has changed between 2013 and 2019 because the Habitat for the species trend has changed from stable to unknown [note that the reason for change is due to less information/accuracy or certainty in the information available].

#### 12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)

a) Unit

number of map 1x1 km grid cells (grids1x1)

- b) Minimum
- c) Maximum
- d) Best single value 603

12.2 Type of estimate

Minimum

12.3 Population size inside the network Method used

Based mainly on extrapolation from a limited amount of data

12.4 Short-term trend of population size within the network Direction

Unknown (x)

12.5 Short-term trend of population size within the network Method used

Based mainly on extrapolation from a limited amount of data

12.6 Additional information

### 13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

### Distribution Map




Figure 1: UK distribution map for S1096 - Brook lamprey (*Lampetra planeri*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available species records within the current reporting period. For further details see the 2019 Article 17 UK Approach document.

### Range Map




Figure 2: UK range map for S1096 - Brook lamprey (*Lampetra planeri*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by applying a bespoke range mapping tool for Article 17 reporting (produced by JNCC) to the 10km grid square distribution map presented in Figure 1. The alpha value for this species was 25km. For further details see the 2019 Article 17 UK Approach document.