European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Conservation status assessment for the species:

S1304 - Greater horseshoe bat (*Rhinolophus* ferrumequinum)

UNITED KINGDOM

IMPORTANT NOTE - PLEASE READ

- The information in this document represents the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- It is based on supporting information provided by the geographically-relevant Statutory Nature Conservation Bodies, which is documented separately.
- The 2019 Article 17 UK Approach document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Maps showing the distribution and range of the species are included (where available).
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the UK assessments. Further underpinning explanatory notes are available in the related country-level reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species).
- The UK-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

NATIONAL LEVEL		
1. General information		
1.1 Member State	UK	
1.2 Species code	1304	
1.3 Species scientific name	Rhinolophus ferrumequinum	
1.4 Alternative species scientific name		
1.5 Common name (in national language)	Greater horseshoe bat	

2. Maps

2.1 Sensitive species	No
2.2 Year or period	1995-2016
2.3 Distribution map	Yes
2.4 Distribution map Method used	Complete survey or a statistically robust estimate
2.5 Additional maps	No

3. Information related to	Annex V Species (Art. 14)	
3.1 Is the species taken in the wild/exploited?	No	
3.2 Which of the measures in Art. 14 have been taken?	a) regulations regarding access to property	No
	b) temporary or local prohibition of the taking of specimens in the wild and exploitation	No
	c) regulation of the periods and/or methods of taking specimens	No
	d) application of hunting and fishing rules which take account of the conservation of such populations	No
	e) establishment of a system of licences for taking specimens or of quotas	No
	f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens	No
	g) breeding in captivity of animal species as well as artificial propagation of plant species	No

h) other measures

No

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

b) Statistics/ quantity taken	Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period					
	Season/	Season/	Season/	Season/	Season/	Season/
	year 1	year 2	year 3	year 4	year 5	year 6
Min. (raw, ie.						
not rounded)						
Max. (raw, ie.						
not rounded)						
Unknown	No	No	No	No	No	No

3.4. Hunting bag or quantity taken in the wild Method used

3.5. Additional information

BIOGEOGRAPHICAL LEVEL

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

4.2 Sources of information

Atlantic (ATL)

England

Arnold, H., 1993. Atlas of Mammals in Britain. (Institute of Terrestrial Ecology Research Publication no. 6), London.

Barrett-Hamilton, G.E., Hinton, M., 1910-1921. A history of British mammals. Gurney and Jackson, London.

Bat Conservation Trust. 2017. The National Bat Monitoring Programme. Annual Report 2016. Bat Conservation Trust, London. Available at

(www.bats.org.uk/pages/nbmp annual report.html)

Bat Conservation Trust. 2018. The National Bat Monitoring Programme. Annual Report 2017. Bat Conservation Trust, London. Available at

http://www.bats.org.uk/pages/results_and_reports.html

Harris, S., Morris, P., Wray, S., Yalden, D., 1995. A review of British mammals: population estimates and conservation status of British Mammals other than cetaceans. JNCC, Peterborough.

JNCC., 2013. Third Report by the United Kingdom under Article 17 on the implementation of the Habitats Directive from January 2007 to December 2012. Species S1304 - Greater horseshoe bat (Rhinolophus ferrumequinum),

Peterborough: JNCC, Available from: www.jncc.gov.uk/article17.

Mathews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C., McDonald, R.A., Shore, R.F., 2018. A review of the population and conservation status of British Mammals. A report by The Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage.

Millais, J., Rothschild, C., Wheler, E., 1904. The mammals of Great Britain and

Ireland. Longmans Green & Co, London. Ransome, R., 1989. Population changes of greater horseshoe bats studied near

Bristol over the past twenty-six years. Biological Journal of the Linnean Society 38, 71-82.

Ransome, R., 1990. The Natural History of Hibernating Bats. Christopher Helm Mammal Series, London.

Ransome, R., 1997. The management of greater horseshoe bat feeding areas to enhance population levels. English Nature Research Reports 241, Peterborough. Ransome, R., 2000. Monitoring diets and population changes of greater horseshoe bats in Gloucestershire and Somerset. English Nature Research Report 341, Peterborough.

Ransome R., Jones, G. 2008. Greater horseshoe bat (Rhinolophus ferrumequinum). Pp. 298-306 in Harris, S & Yalden, D.W. Mammals of the British Isles: Handbook, 4th edition. The Mammal Society, Southampton.

Ransome, R., McOwat, T., 1994. Birth timing and population changes in greater horseshoe bat colonies (Rhinolophus ferrumequinum) are synchronized by climatic temperature. Zoological Journal of the Linnean Society 112, 337-351. Stebbings, R.E., 1988. The conservation of European bats. Christopher Helm Publishers Limited.

Wales

Andrews PT. 2011. Monitoring of Horseshoe Bats through the use of automatic bat counters. Report on nursery roosts in Wales, 2009. Unpublished Report to CCW. CCW, Bangor.

Barr CJ, Gillespie MK. 2000. Estimating hedgerow length and pattern characteristics in Great Britain using Countryside Survey data. Journal of Environmental Management, 60, 23-32.

Bat Conservation Trust. 2018. The State of the UK's Bats 2017. Bat Conservation Trust, London. Available at

(http://www.bats.org.uk/pages/results_and_reports.html)

Bat Conservation Trust. 2018a. The National Bat Monitoring Programme. Annual Report 2017. Bat Conservation Trust, London. Available at

(www.bats.org.uk/pages/nbmp_annual_report.html)

Bat Conservation Trust. 2017. The National Bat Monitoring Programme. Raw Data provided to NRW. Bat Conservation Trust, London

Battersby J. (Ed.). 2005. UK Mammals: Species Status and Population Trends.

JNCC/Tracking Mammals Partnership. JNCC, Peterborough Billington G, Rawlinson MD. 2006. Report on horseshoe bat flight lines and

feeding areas. CCW Science Report No. 775. CCW, Bangor. Dietz C, Helversen OV, Nill D. 2009. Bats of Britain, Europe & Northwest Africa. A

& C Black Publishers Ltd., London.

Dietz C, Keifer A. 2016. Bats of Britain and Europe. London, Bloomsbury Duverge

PL, Jones G. 1994. Greater Horseshoe Bats - activity foraging behaviour and habitat use. British Wildlife 6: 69-77.

English Nature. 2000. Managing landscapes for the greater horseshoe bat. English Nature, Peterborough.

Fensome AG, Mathews F. 2016. Roads and bats: a meta-analysis and review of the evidence on vehicle collisions and barrier effects. Mammal Review, 46(4), 311-323.

Flanders J, Jones G. 2009. Roost use, ranging behaviour and diet of the greater horseshoe bat using non-breeding roosts in Dorset. Journal of Mammalogy 90 888-896

Fox R, Conrad KF, Parsons MF, Warren MS, Woiwod IP. 2006. The state of Britain's larger moths. Butterfly Conservation and Rothamsted Research, Wareham, Dorset.

Harris S, Morris P, Wray S, Yalden D. 1995. A review of British Mammals:

Peterborough.

population estimates and conservation status of British mammals other than cetaceans. JNCC, Peterborough.

Mathews F, Kubasiewicz LM, Gurnell J, Harrower C, McDonald RA, Shore RF. 2018. A review of the population and conservation status of British Mammals. A report by The Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage.

McCracken DI. 1993. The potential for avermectins to affect wildlife. Vet Parasitol, 48(1-4), 273-280.

Natural Resources Wales, 2013. Supporting documentation for the Third Report by the United Kingdom under Article 17 on the implementation of the Directive from January 2007 to December 2012. Conservation status assessment for Species: S1304 - Greater horseshoe bat (Rhinolophus ferrumequinum). Priddis D, Ransome RD & Matthews JE. 2007. Long-distance horseshoes. Bat News, Issue 83, Summer 2007. Bat Conservation Trust, London. Ransome RD. 1989. Population changes of greater horseshoe bats studied near Bristol over the past 26 years. Biological Journal of the Linnean Society, 38, 71-82 Ransome RD. 1990. The Natural History of Hibernating Bats. Christopher Helm. Ransome RD. 1996. The management of feeding areas for greater horseshoe

bats. English Nature Research Report No. 174, English Nature: 1 - 74
Ransome RD. 1997. The management of greater horseshoe bat feeding areas to enhance population levels. English Nature Research Reports 241. Peterborough.
Ransome RD. 1998. The impact of maternity roost conditions on populations of greater horseshoe bats. English Nature Research Report No. 292, English Nature.
Ransome RD. 2000. Monitoring diets and population changes of greater horseshoe bats in Gloucestershire and Somerset. English Nature Report 341.

Ransome RD, Jones G. 2008. Greater horseshoe bat (Rhinolophus ferrumequinum). Pp. 298-306 in Harris, S & Yalden, D.W. Mammals of the British Isles: Handbook, 4th edition. The Mammal Society, Southampton.799pp. Ransome, RD McOwat TP. 1994. Birth timing and population changes in greater horseshoe bat colonies (Rhinolophus ferrumequinum) are synchronised by climatic temperature. Zoological Journal of the Linnean Society, 112, 337-351. Richardson P. 2000. Distribution atlas of bats in Britain and Ireland 1980-1999. Bat Conservation Trust, London.

Robinson RA, Learmonth JA, Hutson AM, Macleod CD, Sparks TH, Leech DI, Pierce GJ, Rehfisch MM, Crick HQP. 2005 Climate change and migratory species. BTO, Thetford.

Rossiter SJ, Jones G, Ransome RD, Barratt EM. 2000. Parentage, reproductive success and breeding behaviour in the greater horseshoe bat (Rhinolophus ferrumequinum). Proceedings of the Royal Society of London. Series B: Biological Sciences, 267(1443), 545-551.

Rossiter SJ, Jones G, Ransome RD, Barratt EM. 2001. Outbreeding increases offspring survival in wild greater horseshoe bats (Rhinolophus ferrumequinum). Proc Biol Sci, 268(1471), 1055-1061.

Rossiter S J, Ransome RD, Faulkes CG, Le Comber SC, Jones G. 2005. Mate fidelity and intra-lineage polygyny in greater horseshoe bats. Nature, 437(7057), 408-411.

Russ J. 2012. British bat calls: a guide to species identification. Exeter, Pelagic Publishing.

Spencer JW, Kirby KJ. 1992. An inventory of ancient woodland for England and Wales. Biological Conservation, 62, 77-93

Stebbings RE. 1989. Conservation of the greater horseshoe bat: is the long term survival of the greater horseshoe bat a viable concept? British Wildlife, 1, 14-19.

Stone EL, Jones G, Harris S. 2009. Street Lighting Disturbs Commuting Bats. Current Biology, 19(13), 1123-1127.

5. Range

5.1 Surface area (km²)

43015

5.2 Short-term trend Period

2013-2018

a) Minimum

5.3 Short-term trend Direction

Increasing (+)

5.4 Short-term trend Magnitude

b) Maximum

5.5 Short-term trend Method used

Complete survey or a statistically robust estimate

5.6 Long-term trend Period

5.7 Long-term trend Direction

5.8 Long-term trend Magnitude

a) Minimum

b) Maximum

5.9 Long-term trend Method used

5.10 Favourable reference range

a) Area (km²)

43015

b) Operator

c) Unknown

d) Method

The FRR has changed since 2013. The new value is considered to be large enough to support a viable population and no lower than the range estimate when the Habitats Directive came into force in the UK. For further information see the 2019 Article 17 UK Approach document. The 2013 FRR value has been revised and is equal to the current range. The current range surface area has been calculated using the method outlined in Mathews et. al., (2018) and is based on presence data collected between 1995-2016. Areas that contain very isolated records may not have been included in the area of distribution. The new, more robust method of calculating range has reduced estimated range size for this species since 2013. This does not represent a real reduction in range.

5.11 Change and reason for change in surface area of range

Genuine change Improved knowledge/more accurate data Use of different method

The change is mainly due to: Use of different method

5.12 Additional information

Short term trend in range has been assessed by using the 2019 distribution data and the 2013 method for calculating range, and comparing the result with range surface area in 2013. For further information see the 2019 Article 17 UK Approach document and country assessments.

6. Population

6.1 Year or period

1995-2017

6.2 Population size (in reporting unit) a) Unit

a) Unit number of individuals (i)

b) Minimum 9245c) Maximum 18530

d) Best single value 12951

6.3 Type of estimate

Best estimate

6.4 Additional population size (using population unit other than reporting unit)

a) Unit

b) Minimum

c) Maximum

d) Best single value

6.5 Type of estimate

6.6 Population size Method used

Complete survey or a statistically robust estimate

6.7 Short-term trend Period

2006-2017

6.8 Short-term trend Direction

Increasing (+)

6.9 Short-term trend Magnitude

a) Minimum

b) Maximum

c) Confidence interval

6.10 Short-term trend Method used

Complete survey or a statistically robust estimate

6.11 Long-term trend Period

6.12 Long-term trend Direction

6.13 Long-term trend Magnitude

- a) Minimum
- b) Maximum
- c) Confidence interval

6.14 Long-term trend Method used

6.15 Favourable reference population (using the unit in 6.2 or 6.4)

a) Population size

12951 with unit number of individuals (i)

- b) Operator
- c) Unknown

d) Method

The FRP has changed since 2013. The new value is considered to be large enough to support a viable population and no less than the population estimate when the Habitats Directive came into force in the UK. For further information see the 2019 Article 17 UK Approach document. The FRP is the same as the new population estimate (Mathews et. al., 2018). The previous FRP was thought to be an underestimate of the population, currently, and when the Habitats Directive came into force in the UK.

6.16 Change and reason for change in population size

Genuine change

Improved knowledge/more accurate data

The change is mainly due to: Genuine change

6.17 Additional information

The UK National Bat Monitoring Programme (NBMP) has detected increases in both the maternity and hibernation indices and these changes have been

consistent over this reporting period. For further information see the 2019 Article 17 UK Approach document.

7. Habitat for the species

7.1 Sufficiency of area and quality of occupied habitat

a) Are area and quality of occupied habitat sufficient (for long-term survival)?

Yes

b) Is there a sufficiently large area of unoccupied habitat of suitable quality (for long-term survival)?

7.2 Sufficiency of area and quality of occupied habitat Method used

Based mainly on extrapolation from a limited amount of data

occupied habitat Method used
7.3 Short-term trend Period

1995-2016

7.4 Short-term trend Direction

Stable (0)

7.5 Short-term trend Method used

Based mainly on extrapolation from a limited amount of data

7.6 Long-term trend Period

7.0 Long-term trend remod

7.7 Long-term trend Direction

7.8 Long-term trend Method used

7.9 Additional information

This species uses a mosaic of habitats. Given that the population appears to be increasing and the range is stable, the habitat can also be considered to be stable

8. Main pressures and threats

8.1 Characterisation of pressures/threats

Pressure	Ranking
Removal of small landscape features for agricultural land parcel consolidation (hedges, stone walls, rushes, open ditches, springs, solitary trees, etc.) (A05)	Н
Abandonment of grassland management (e.g. cessation of grazing or mowing) (A06)	Н
Livestock farming (without grazing) (A14)	Н
Logging without replanting or natural regrowth (B05)	M
Extraction of minerals (e.g. rock, metal ores, gravel, sand, shell) (C01)	M
Roads, paths, railroads and related infrastructure (e.g. bridges, viaducts, tunnels) (E01)	Н
Construction or modification (e.g. of housing and settlements in existing urban or recreational areas (FO2)) H
Sports, tourism and leisure activities (F07)	M
Reduced fecundity / genetic depression (e.g. inbreeding or endogamy) (L05)	M
Other natural catastrophes (M10)	M
Threat	Ranking
Removal of small landscape features for agricultural land parcel consolidation (hedges, stone walls, rushes, open	М

ditches, springs, solitary trees, etc.) (A05)	
Abandonment of grassland management (e.g. cessation of grazing or mowing) (A06)	Н
Livestock farming (without grazing) (A14)	M
Logging without replanting or natural regrowth (B05)	M
Extraction of minerals (e.g. rock, metal ores, gravel, sand, shell) (CO1)	M
Roads, paths, railroads and related infrastructure (e.g. bridges, viaducts, tunnels) (E01)	Н
Construction or modification (e.g. of housing and settlements) in existing urban or recreational areas (F02)	Н
Sports, tourism and leisure activities (F07)	M
Reduced fecundity / genetic depression (e.g. inbreeding or endogamy) (L05)	Н
Other natural catastrophes (M10)	Н

8.2 Sources of information

8.3 Additional information

9. Conservation measures

9.1 Status of measures	a) Are measures needed?	Yes
J.I Status of fileasures	a) Alt illeasures liceacu:	103

b) Indicate the status of measures Measures identified and taken

9.2 Main purpose of the measures Maintain the current range, population and/or habitat for the species taken

9.3 Location of the measures taken Both inside and outside Natura 2000

9.4 Response to the measures Medium-term results (within the next two reporting periods, 2019-2030)

9.5 List of main conservation measures

Restore small landscape features on agricultural land (CA02)

Adapt mowing, grazing and other equivalent agricultural activities (CA05)

Manage the use of natural fertilisers and chemicals in agricultural (plant and animal) production (CA09)

Adapt/change forest management and exploitation practices (CB05)

Reduce impact of transport operation and infrastructure (CE01)

Manage/reduce/eliminate noise, light and other forms of pollution from transport (CE05)

Manage conversion of land for construction and development of infrastructure (CF01)

Reduce impact of outdoor sports, leisure and recreational activities (CF03)

Improvement of habitat of species from the directives (CS03)

9.6 Additional information

10. Future prospects

10.1 Future prospects of parameters

a) Range Good b) Population Good c) Habitat of the species Good

10.2 Additional information

Future trend in Range is Positive - increasing <=1% (one percent or less) per year on average; Future trend in Population is Very Positive - increasing >1% (more than one percent) per year on average; and Future trend in Habitat for the species is Overall stable. For further information on how future trends inform the Future Prospects conclusion see the 2019 Article 17 UK Approach document.

11. Conclusions

11.3. Habitat for the species

11.5 Overall assessment of

11.4. Future prospects

Conservation Status

11.1. Range Favourable (FV)

11.2. Population Favourable (FV)

Favourable (FV)

Favourable (FV)

Favourable (FV)

11.6 Overall trend in Conservation

Improving (+)

Status

a) Overall assessment of conservation status

11.7 Change and reasons for change in conservation status and conservation status trend

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

11.8 Additional information

Conclusion on Range reached because: (i) the short-term trend direction in Range surface area is increasing; and (ii) the current Range surface area is approximately equal to the Favourable Reference Range.

Conclusion on Population reached because: (i) the short-term trend direction in Population size is increasing; and (ii) the current Population size is approximately equal to the Favourable Reference Population.

Conclusion on Habitat for the species reached because: (i) the area of [occupied habitat is [sufficiently large and (ii) the habitat quality is suitable for the long-term survival of the species; and (iii) the short-term trend in area and quality of habitat is stable.

Conclusion on Future prospects reached because: (i) the Future prospects for Range are good; (ii) the Future prospects for Population good; and (iii) the Future prospects for Habitat for the species are good.

Overall assessment of Conservation Status is Favourable because all of the conclusions are Favourable.

Overall trend in Conservation Status is based on the combination of the short-term trends for Range - increasing, Population - increasing, and Habitat for the species - stable.

Overall assessment of Conservation Status has not changed since 2013. The overall trend in conservation status was not reported for this species in 2013. However, from the information available the overall trend would have been increasing in 2013 and so there has been no change since the last reporting

round.

12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)

a) Unit number of individuals (i)

- b) Minimum
- c) Maximum
- d) Best single value 4734

12.2 Type of estimate

12.3 Population size inside the network Method used

Minimum

Complete survey or a statistically robust estimate

12.4 Short-term trend of population size within the network Direction

Increasing (+)

12.5 Short-term trend of population size within the network Method used

Complete survey or a statistically robust estimate

12.6 Additional information

13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

Distribution Map




Figure 1: UK distribution map for S1304 - Greater horseshoe bat (*Rhinolophus ferrumequinum*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available species records within the current reporting period. For further details see the 2019 Article 17 UK Approach document.

Range Map




Figure 2: UK range map for S1304 - Greater horseshoe bat (*Rhinolophus ferrumequinum*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by The Mammal Society applying a range mapping tool as outlined in Matthews et al. (2018), to the 10km grid square distribution map presented in Figure 1. The alpha value for this species was 20km. For further details see the 2019 Article 17 UK Approach document.