

**European Community Directive
on the Conservation of Natural Habitats
and of Wild Fauna and Flora
(92/43/EEC)**

**Fourth Report by the United Kingdom
under Article 17**

on the implementation of the Directive
from January 2013 to December 2018

Conservation status assessment for the species:

S1309 - Common pipistrelle (*Pipistrellus pipistrellus*)

UNITED KINGDOM

IMPORTANT NOTE - PLEASE READ

- The information in this document represents the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- It is based on supporting information provided by the geographically-relevant Statutory Nature Conservation Bodies, which is documented separately.
- The 2019 Article 17 UK Approach document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Maps showing the distribution and range of the species are included (where available).
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the UK assessments. Further underpinning explanatory notes are available in the related country-level reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species).
- The UK-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, <https://jncc.gov.uk/article17>, for further information on UK Article 17 reporting.

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

NATIONAL LEVEL

1. General information

1.1 Member State	UK
1.2 Species code	1309
1.3 Species scientific name	<i>Pipistrellus pipistrellus</i>
1.4 Alternative species scientific name	
1.5 Common name (in national language)	Common pipistrelle

2. Maps

2.1 Sensitive species	No
2.2 Year or period	1994-2018
2.3 Distribution map	Yes
2.4 Distribution map Method used	Complete survey or a statistically robust estimate
2.5 Additional maps	No

3. Information related to Annex V Species (Art. 14)

3.1 Is the species taken in the wild/exploited?	No	
3.2 Which of the measures in Art. 14 have been taken?	a) regulations regarding access to property	No
	b) temporary or local prohibition of the taking of specimens in the wild and exploitation	No
	c) regulation of the periods and/or methods of taking specimens	No
	d) application of hunting and fishing rules which take account of the conservation of such populations	No
	e) establishment of a system of licences for taking specimens or of quotas	No
	f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens	No
	g) breeding in captivity of animal species as well as artificial propagation of plant species	No
	h) other measures	No

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

b) Statistics/ quantity taken	Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period					
	Season/ year 1	Season/ year 2	Season/ year 3	Season/ year 4	Season/ year 5	Season/ year 6
Min. (raw, ie. not rounded)						
Max. (raw, ie. not rounded)						
Unknown	No	No	No	No	No	No

3.4. Hunting bag or quantity taken in the wild Method used

3.5. Additional information

BIOGEOGRAPHICAL LEVEL

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

Atlantic (ATL)

4.2 Sources of information

England

Barratt, E.M., R. Deaville, T.M. Burland, M.W. Bruford, G. Jones, P.A. Racey and R.K. Wayne (1997). DNA answers the call of pipistrelle bat species. *Nature*, 387 (6629), 138-139

Bat Conservation Trust (2018). The State of the UK's Bats 2017. Bat Conservation Trust, London. Available at (http://www.bats.org.uk/pages/results_and_reports.html)

Boye, P and Dietz, M. (2005). Research Report No 661: Development of good practice guidelines for woodland management for bats. English Nature, Peterborough.

Davidson-Watts, I. and Jones, G. (2006). Differences in foraging behaviour between *Pipistrellus pipistrellus* (Schreber, 1774) and *Pipistrellus pygmaeus* (Leach, 1825). *Journal of Zoology*, 268 (1), 55-62

Fensome, A. G. and Mathews, F. (2016). Roads and bats: a meta-analysis and review of evidence on vehicle collisions and barrier effects. *Mammal Review*, 46 (4), 311-323

Fuentes-Montemayor, E., Goulson, D., Cavin, L., Wallace, J.M., and Park, K.J. (2013). Fragmented woodlands in agricultural landscapes: The influence of woodland character and landscape context on bats and their insect prey. *Agriculture, Ecosystems and Environment*, 172, 6-15

Glendell, M. and Vaughan, N. (2002). Foraging activity of bats in historic landscapoe parks in relation to habitat composition and park management. *Animal Conservation*, 5 (4), 309-316

Jones, G and Racey, P.A. (2008). Common pipistrelle *Pipistrellus pipistrellus*, Soprano pipistrelle *Pipistrellus pygmaeus*. Pages 343-351 In Harris, S and Yalden,

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

- D.W. Mammals of the British Isles: Handbook, 4th edition. The Mammal Society, Southampton. 799pp.
- Lintott, P.R., Bunnefeld, N. and Park, K.J. (2015). Opportunities for improving the foraging potential of urban waterways for bats. *Biological Conservation*, 191, 224-233.
- Lintott, P.R., Barlow, K., Bunnefeld, N., Briggs, P., Gajas Roig, C., and Park, K.J. (2016). Differential responses of cryptic bat species to the urban landscape. *Ecology and Evolution*, 6 (7), 2044-2052
- Mathews, F., Richardson, S.M., and Hosken, D.J. (2016). Understanding the risks to bat populations posed by wind turbines - Phase 2 - WC0753, Defra.
- Mathews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C., McDonald, R.A., Shore, R.F (2018). A review of the population and conservation status of British Mammals. A report by the Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage.
- Mitchell-Jones, T.J. (2010). Bats in houses - the conservation challenge. Pp 3965-378 in *Species Management : challenges and solutions for the 21st century*.
- Baxter, J.M. and Galbraith, C.A. TSO Scotland, Edinburgh
- Nicholls, B. and Racey, P (2006a). Habitat selection as a mechanism of resource partitioning in two cryptic bat species *Pipistrellus pipistrellus* and *pipistrellus pygmaeus*. *Ecography*, 29, 697-708
- Nicholls, B. and Racey, P (2006b). Contrasting home-range size and spatial partitioning in cryptic and sympatric bats. *Behavioural Ecology and Sociobiology*, 61, 131-142
- Waring, S.D., Essah, E., Gunnell, K, and Bonser, R (2013). Double jeopardy: the potential for problems when bats interact with breathable roofing membranes in the United Kingdom. *Architecture and Environment*, 1 1-3
- Warren, R, D., Waters, D, A., Altringham, J.D., and Bullock, D.J. (2000). The distribution of Daubenton's bats (*Myotis daubentonii*) and pipistrelle bats (*Pipistrellus pipistrellus*) (Vespertilionidae) in relation to small-scale variation in riverine habitat. *Biological Conservation*, 92 (1), 85-91
- Scotland
- Barratt, E.M., R. Deaville, T.M. Burland, M.W. Bruford, G. Jones, P.A. Racey and R.K. Wayne (1997). DNA answers the call of pipistrelle bat species. *Nature*, 387 (6629), 138-139
- Bat Conservation Trust (2018). The State of the UK's Bats 2017. Bat Conservation Trust, London. Available at (http://www.bats.org.uk/pages/results_and_reports.html)
- Boye, P and Dietz, M. (2005). Research Report No 661: Development of good practice guidelines for woodland management for bats. English Nature, Peterborough.
- Davidson-Watts, I. and Jones, G. (2006). Differences in foraging behaviour between *Pipistrellus pipistrellus* (Schreber, 1774) and *Pipistrellus pygmaeus* (Leach, 1825). *Journal of Zoology*, 268 (1), 55-62
- Fensome, A. G. and Mathews, F. (2016). Roads and bats: a meta-analysis and review of evidence on vehicle collisions and barrier effects. *Mammal Review*, 46 (4), 311-323
- Fuentes-Montemayor, E., Goulson, D., Cavin, L., Wallace, J.M., and Park, K.J. (2013). Fragmented woodlands in agricultural landscapes: The influence of woodland character and landscape context on bats and their insect prey. *Agriculture, Ecosystems and Environment*, 172, 6-15
- Glendell, M. and Vaughan, N. (2002). Foraging activity of bats in historic landscape parks in relation to habitat composition and park management. *Animal Conservation*, 5 (4), 309-316

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

- Jones, G and Racey, P.A. (2008). Common pipistrelle *Pipistrellus pipistrellus*, Soprano pipistrelle *Pipistrellus pygmaeus*. Pages 343-351 In Harris, S and Yalden, D.W. Mammals of the British Isles: Handbook, 4th edition. The Mammal Society, Southampton. 799pp.
- Lintott, P.R., Bunnefeld, N. and Park, K.J. (2015). Opportunities for improving the foraging potential of urban waterways for bats. *Biological Conservation*, 191, 224-233.
- Lintott, P.R., Barlow, K., Bunnefeld, N., Briggs, P., Gajas Roig, C., and Park, K.J. (2016). Differential responses of cryptic bat species to the urban landscape. *Ecology and Evolution*, 6 (7), 2044-2052
- Mathews, F., Richardson, S.M., and Hosken, D.J. (2016). Understanding the risks to bat populations posed by wind turbines - Phase 2 - WC0753, Defra.
- Mathews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C., McDonald, R.A., Shore, R.F (2018). A review of the population and conservation status of British Mammals. A report by the Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage.
- Mitchell-Jones, T.J. (2010). Bats in houses - the conservation challenge. Pp 3965-378 in *Species Management : challenges and solutions for the 21st century*.
- Baxter, J.M. and Galbraith, C.A. TSO Scotland, Edinburgh
- Newson, S.E., Evans, H.E., Gillings, S., Jarrett, D. & Wilson, M.W. 2017. A survey of high risk bat species across southern Scotland. Scottish Natural Heritage Commissioned Report No. 1008.
- Nicholls, B. and Racey, P (2006a). Habitat selection as a mechanism of resource partitioning in two cryptic bat species *Pipistrellus pipistrellus* and *pipistrellus pygmaeus*. *Ecography*, 29, 697-708
- Nicholls, B. and Racey, P (2006b). Contrasting home-range size and spatial partitioning in cryptic and sympatric bats. *Behavioural Ecology and Sociobiology*, 61, 131-142
- Waring, S.D., Essah, E., Gunnell, K, and Bonser, R (2013). Double jeopardy: the potential for problems when bats interact with breathable roofing membranes in the United Kingdom. *Architecture and Environment*, 1 1-3
- Warren, R, D., Waters, D, A., Altringham, J.D., and Bullock, D.J. (2000). The distribution of Daubenton's bats (*Myotis daubentonii*) and pipistrelle bats (*Pipistrellus pipistrellus*) (Vespertilionidae) in relation to small-scale variation in riverine habitat. *Biological Conservation*, 92 (1), 85-91
- Wales
- Bat Conservation Trust. 2018. The State of the UK's Bats 2017. Bat Conservation Trust, London. Available at http://www.bats.org.uk/pages/results_and_reports.html
- Bat Conservation Trust. 2018a. The National Bat Monitoring Programme. Annual Report 2017. Bat Conservation Trust, London. Available at www.bats.org.uk/pages/nbmp_annual_report.html
- Bat Conservation Trust. 2017. The National Bat Monitoring Programme. Raw Data provided to NRW. Bat Conservation Trust, London
- Barlow KE. 1997. The diets of two phonic types of the bat *Pipistrellus pipistrellus* in Britain. *Journal of Zoology*, 243(3), 597-609.
- Barlow KE, Jones G. 1999. Roosts, echolocation calls and wing morphology of two phonic types of *Pipistrellus pipistrellus*. *Zeitschrift fur Saugetierkunde*, 64, 257-268.
- Barratt EM, Deaville R, Burland TM, Bruford MW, Jones G, Racey PA, Wayne RK. 1997. DNA answers the call of pipistrelle bat species. *Nature (Lond.)*, 387:138-139.
- Battersby J. (Ed.). 2005. UK Mammals: Species Status and Population Trends.

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

- JNCC/Tracking Mammals Partnership. JNCC, Peterborough
- Boye P, Dietz M. 2005. Research Report No 661: Development of good practice guidelines for woodland management for bats. English Nature, Peterborough.
- Davidson-Watts I, Jones G. 2006. Differences in foraging behaviour between *Pipistrellus pipistrellus* (Schreber, 1774) and *Pipistrellus pygmaeus* (Leach, 1825). *Journal of Zoology*, 268: 55-62.
- Dietz C, Helversen OV, Nill D. 2009. Bats of Britain, Europe & Northwest Africa. A & C Black Publishers Ltd., London.
- Dietz C, Keifer A. 2016. Bats of Britain and Europe. London, Bloomsbury
- Fensome AG, Mathews F. 2016. Roads and bats: a meta-analysis and review of evidence on vehicle collisions and barrier effects. *Mammal Review*, 46 (4), 311-323
- Fuentes-Montemayor E, Goulson D, Cavin L, Wallace JM, Park KJ. 2013. Fragmented woodlands in agricultural landscapes: The influence of woodland character and landscape context on bats and their insect prey. *Agriculture, Ecosystems and Environment*, 172, 6-15
- Glendell M. Vaughan N. 2002. Foraging activity of bats in historic landscape parks in relation to habitat composition and park management. *Animal Conservation*, 5 (4), 309-316
- Harris S, Morris P, Wray S, Yalden D. 1995. A review of British Mammals: population estimates and conservation status of British mammals other than cetaceans. JNCC, Peterborough.
- Jenkins EV, Laine T, Morgan SE, Cole KR, Speakman JR. 1998. Roost selection in the pipistrelle bat, *Pipistrellus pipistrellus* (Chiroptera: Vespertilionidae), in northeast Scotland. *Anim Behav*, 56(4), 909-917.
- Jones G, Barratt EM. 1999. *Vespertilio pipistrellus* Schreber, 1774 and *V. pygmaeus* Leach, 1825 (currently *Pipistrellus pipistrellus* and *P. pygmaeus*; Mammalia, Chiroptera): proposed designation of neotypes. *Bulletin of Zoological Nomenclature*, 56: 182-186.
- Jones G, Racey PA. 2008. Common pipistrelle *Pipistrellus pipistrellus*, Soprano pipistrelle *Pipistrellus pygmaeus*. Pages 343-351 In Harris, S. & Yalden, D.W. *Mammals of the British Isles: Handbook*, 4th edition. The Mammal Society, Southampton. 799pp.
- Lintott PR, Bunnefeld N, Park KJ. 2015. Opportunities for improving the foraging potential of urban waterways for bats. *Biological Conservation*, 191, 224-233.
- Lintott PR, Barlow K, Bunnefeld N, Briggs P, Gajas Roig C, Park KJ. 2016. Differential responses of cryptic bat species to the urban landscape. *Ecology and Evolution*, 6 (7), 2044-2052
- Mathews F, Richardson SM, Hosken DJ. 2016. Understanding the risks to bat populations posed by wind turbines - Phase 2 - WC0753, Defra.
- Mathews F, Kubasiewicz LM, Gurnell J, Harrower C, McDonald RA, Shore RF. 2018. A review of the population and conservation status of British Mammals. A report by The Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage.
- Mitchell-Jones TJ. 2010. Bats in houses - the conservation challenge. Pp 365-378, In: *Species Management: challenges and solutions for the 21st century*. Baxter, J.M. & Galbraith, C.A. Tso Scotland, Edinburgh.
- Mitchell-Jones TJ, Carlin C. 2009. TIN051 Bats and onshore wind turbines Interim Guidance. 2nd edition, February 2012. Available at <http://publications.naturalengland.org.uk/file/490077>
- Natural Resources Wales, 2013. Supporting documentation for the Third Report by the United Kingdom under Article 17 on the implementation of the Directive from January 2007 to December 2012. Conservation status assessment for

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

- Species: S1309 - Common pipistrelle bat (*Pipistrellus pipistrellus*)
- Nicholls B, Racey P. 2006a. Habitat selection as a mechanism of resource partitioning in two cryptic bat species *Pipistrellus pipistrellus* and *Pipistrellus pygmaeus*. *Ecography*, 29, 697-708.
- Nicholls B, Racey P. 2006b. Contrasting home-range size and spatial partitioning in cryptic and sympatric pipistrelle bats. *Behavioural Ecology and Sociobiology*, 61, 131-142.
- Oakley SF, Jones G. 1998. Habitat around maternity roosts of the 55 kHz phonic type of pipistrelle bats (*Pipistrellus pipistrellus*). *Journal of Zoology*, 245(2), 222-228.
- Racey PA. 1969. Diagnosis of pregnancy and experimental extension of gestation in the pipistrelle bat, *Pipistrellus pipistrellus*. *J Reprod Fertil*, 19(3), 465-474.
- Richardson P. 2000. Distribution atlas of bats in Britain and Ireland 1980-1999. Bat Conservation Trust, London.
- Speakman JR. 1991. The impact of predation by birds on bat populations in the British Isles. *Mammal Review*, 21, 123-142.
- Waring SD, Essah E, Gunnell K, Bonser R. 2013. Double jeopardy: the potential for problems when bats interact with breathable roofing membranes in the United Kingdom. *Architecture and Environment*, 1 1-3
- Warren RD, Waters DA, Altringham JD, Bullock DJ. 2000. The distribution of Daubenton's bats (*Myotis daubentonii*) and pipistrelle bats (*Pipistrellus pipistrellus*) (*Vespertilionidae*) in relation to small-scale variation in riverine habitat. *Biological Conservation*, 92 (1), 85-91
- N.Ireland
- Lundy, M. & Montgomery, I. (2010) Summer habitat associations of bats between riparian landscapes and within riparian areas, *European Journal of Wildlife Research*, 56(3): 385-394.
- Lundy, M.G., Aughney, T., Montgomery, W.I., and Roche, N. (2011). Landscape conservation for Irish bats & species: specific roosting characteristics. Bat Conservation Ireland. Unpublished.
- Russ, J.M. & Montgomery, W.I. (2002). Habitat association of bats in Northern Ireland: implications for conservation. *Biological Conservation*. 108: 49-58.
- Lundy, M.G., Buckley, D.J., Boston, E.S.M., Scott, D.D., Prodohl, P.A., Marnell, F., Teeling, E.C., Montgomery, W.I., (2012). Behavioural context of multi-scale species distribution models assessed by radio-tracking. *Basic Appl. Ecol.*, <http://dx.doi.org/10.1016/j.baae.2011.1012.1003>.
- Hutson, A.M., Mickleburgh, S.P., and Racey, P.A. (comp.). (2001). Microchiropteran bats: global status survey and conservation action plan. IUCN/SSC Chiroptera Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK. x + 258 pp.
- Hutson, A.M., Mickleburgh, S.P. & Racey, P.A. (comp.). (2001) Global Status Survey and Conservation Action Plan Microchiropteran Bats, The Nature Conservation Bureau Ltd, ISBN: 2-8317-0595-9, <http://www.uni-giessen.de/faculties/f08/departments/tsz/mammalian-ecology-group/downloads/iucn-microchiroptera>
- Russ, J.M. (1999). The Microchiroptera of Northern Ireland: community composition, habitat associations and ultrasound. Unpublished Ph.D thesis. The Queen's University of Belfast.
- Russ, J.M., Briffa M. & Montgomery, W.I. (2003). Seasonal patterns in activity and habitat use by *Pipistrellus* spp. and *Nyctalus leisleri* in Northern Ireland, determined using a driving transect. *Journal of Zoology*. 259: 289-299.
- Boston, E. (2016) A report on Article 17 reporting for Northern Ireland on the eight bat species listed in annex IV of the UK Habitats Directive, unpublished

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

report compiled for CEDaR

Mathews, F., Richardson, S., Lintott, P., and Hosken, D. 2016. Understanding the Risk to European Protected Species (bats) at Onshore Wind Turbine Sites to inform Risk Management. University of Exeter. Report to DEFRA.

Roche, N., Langton, S. and Aughney T. (2012) Car-based bat monitoring in Ireland 2003-2011. Irish Wildlife Manuals, No. 60. National Parks and Wildlife Service, Department of the Arts, Heritage and the Gaeltacht, Ireland.

Aughney, T., Roche, N., & Langton, S. (2016) Irish Bat Monitoring Schemes: Annual Report for 2015. www.batconservationireland.org.

5. Range

5.1 Surface area (km ²)	233480
5.2 Short-term trend Period	2013-2018
5.3 Short-term trend Direction	Stable (0)
5.4 Short-term trend Magnitude	a) Minimum b) Maximum
5.5 Short-term trend Method used	Complete survey or a statistically robust estimate
5.6 Long-term trend Period	
5.7 Long-term trend Direction	
5.8 Long-term trend Magnitude	a) Minimum b) Maximum
5.9 Long-term trend Method used	
5.10 Favourable reference range	<p>a) Area (km²) 230973</p> <p>b) Operator</p> <p>c) Unknown</p> <p>d) Method</p> <p>The FRR is the same as in 2013. The value is considered to be large enough to support a viable population and no lower than the range estimate when the Habitats Directive came into force in the UK. For further information see the 2019 Article 17 UK Approach document.</p> <p>The current range surface area has been calculated using the method outlined in Mathews et al. (2018) and is based on presence data collected between 1995-2016. Areas that contain very isolated records may not have been included in the area of distribution. The new, more robust method of calculating range has reduced the estimated range size for this species since 2013. This does not represent a real reduction in range. Current range is above the FRR.</p>
5.11 Change and reason for change in surface area of range	<p>Improved knowledge/more accurate data</p> <p>Use of different method</p> <p>The change is mainly due to: Use of different method</p>
5.12 Additional information	Trend in range has been assessed by using the 2019 distribution data and the 2013 method for calculating range, and comparing that with range surface area in 2013. For further details please see the 2019 Article 17 UK Approach document and country assessments.

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

6. Population

6.1 Year or period	1994-2018
6.2 Population size (in reporting unit)	a) Unit number of map 1x1 km grid cells (grids1x1) b) Minimum c) Maximum d) Best single value 31559
6.3 Type of estimate	Minimum
6.4 Additional population size (using population unit other than reporting unit)	a) Unit number of individuals (i) b) Minimum 1100600 c) Maximum 7843000 d) Best single value
6.5 Type of estimate	95% confidence interval
6.6 Population size Method used	Complete survey or a statistically robust estimate
6.7 Short-term trend Period	2003-2018
6.8 Short-term trend Direction	Increasing (+)
6.9 Short-term trend Magnitude	a) Minimum b) Maximum c) Confidence interval
6.10 Short-term trend Method used	Based mainly on extrapolation from a limited amount of data
6.11 Long-term trend Period	
6.12 Long-term trend Direction	
6.13 Long-term trend Magnitude	a) Minimum b) Maximum c) Confidence interval
6.14 Long-term trend Method used	
6.15 Favourable reference population (using the unit in 6.2 or 6.4)	a) Population size b) Operator Approximately equal to (\approx) c) Unknown d) Method <p>The FRP has changed since 2013. An FRP operator has been used because it has not been possible to calculate the exact FRP value. The current population (individuals) is considered to be a viable population and is no less than when the Habitats Directive came into force in the UK. For further details see the 2019 Article 17 UK Approach document. The confidence limits for the population estimate are extremely wide and methodologies have changed. A best single value for the population has not been provided because of the level of uncertainty around the population estimate. Instead the lower and upper confidence intervals provide minimum and maximum limits to the</p>

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

estimate.

6.16 Change and reason for change in population size

Genuine change
Improved knowledge/more accurate data
Use of different method
The change is mainly due to: Genuine change

6.17 Additional information

The 1km square count has been calculated from the UK count of 1km squares where the species has been recorded. This is a minimum count because it only includes number of recorded occupied 1km squares. UK population estimates have been derived from the GB estimate in Mathews et. al., 2018, and the Northern Ireland estimate from the 2013 Article 17 report. The UK National Bat Monitoring Programme (NBMP) has recorded a statistically significant increase in *P. pipistrellus* between 2006-2017. Knowledge of the species has improved. The change is also due to a different method for calculating population size. The current population (in individuals) is considered to be approximately equal to the FRP and is sufficient to maintain a viable population.

7. Habitat for the species

7.1 Sufficiency of area and quality of occupied habitat

a) Are area and quality of occupied habitat sufficient (for long-term survival)? Yes
b) Is there a sufficiently large area of unoccupied habitat of suitable quality (for long-term survival)?

7.2 Sufficiency of area and quality of occupied habitat Method used

Based mainly on extrapolation from a limited amount of data

7.3 Short-term trend Period

2007-2018

7.4 Short-term trend Direction

Stable (0)

7.5 Short-term trend Method used

Based mainly on extrapolation from a limited amount of data

7.6 Long-term trend Period

7.7 Long-term trend Direction

7.8 Long-term trend Method used

7.9 Additional information

The species requires a complex mosaic of habitats to support foraging, roosting and commuting behaviour. The stable habitat trend is based on limited data and expert opinion.

8. Main pressures and threats

8.1 Characterisation of pressures/threats

Pressure	Ranking
Conversion from one type of agricultural land use to another (excluding drainage and burning) (A02)	M
Removal of small landscape features for agricultural land parcel consolidation (hedges, stone walls, rushes, open ditches, springs, solitary trees, etc.) (A05)	M
Use of plant protection chemicals in agriculture (A21)	M

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

Conversion to other types of forests including monocultures (B02)	M
Logging without replanting or natural regrowth (B05)	M
Wind, wave and tidal power, including infrastructure (D01)	H
Roads, paths, railroads and related infrastructure (e.g. bridges, viaducts, tunnels) (E01)	H
Construction or modification (e.g. of housing and settlements) in existing urban or recreational areas (F02)	H
Residential or recreational activities and structures generating noise, light, heat or other forms of pollution (F24)	H
Industrial or commercial activities and structures generating noise, light, heat or other forms of pollution (F25)	H
Threat	Ranking
Conversion from one type of agricultural land use to another (excluding drainage and burning) (A02)	M
Removal of small landscape features for agricultural land parcel consolidation (hedges, stone walls, rushes, open ditches, springs, solitary trees, etc.) (A05)	M
Use of plant protection chemicals in agriculture (A21)	M
Conversion to other types of forests including monocultures (B02)	M
Logging without replanting or natural regrowth (B05)	M
Wind, wave and tidal power, including infrastructure (D01)	M
Roads, paths, railroads and related infrastructure (e.g. bridges, viaducts, tunnels) (E01)	H
Construction or modification (e.g. of housing and settlements) in existing urban or recreational areas (F02)	H
Residential or recreational activities and structures generating noise, light, heat or other forms of pollution (F24)	H
Industrial or commercial activities and structures generating noise, light, heat or other forms of pollution (F25)	H

8.2 Sources of information

8.3 Additional information

9. Conservation measures

9.1 Status of measures

- a) Are measures needed? Yes
- b) Indicate the status of measures Measures identified and taken

9.2 Main purpose of the measures taken

Maintain the current range, population and/or habitat for the species

9.3 Location of the measures taken

Both inside and outside Natura 2000

9.4 Response to the measures

Medium-term results (within the next two reporting periods, 2019-2030)

9.5 List of main conservation measures

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

Other measures related to agricultural practices (CA16)
Prevent conversion of (semi-) natural habitats into forests and of (semi-)natural forests into intensive forest plantation (CB01)
Adapt/manage reforestation and forest regeneration (CB04)
Adapt/manage renewable energy installation, facilities and operation (CC03)
Reduce impact of transport operation and infrastructure (CE01)
Manage conversion of land for construction and development of infrastructure (CF01)
Reduce/eliminate diffuse pollution to surface or ground waters from industrial, commercial, residential and recreational areas and activities (CF05)
Reduce/eliminate noise, light, heat or other forms pollution from industrial, commercial, residential and recreational areas and activities (CF09)
Prevent conversion of natural and semi-natural habitats, and habitats of species into agricultural land (CA01)
Restore small landscape features on agricultural land (CA02)

9.6 Additional information

10. Future prospects

10.1 Future prospects of parameters	a) Range Good b) Population Good c) Habitat of the species Good
10.2 Additional information	Future trend in Range is Overall stable; Future trend in Population is Very Positive - increasing >1% (more than one percent) per year on average; and Future trend in Habitat for the species is Overall stable. For further information on how future trends inform the Future Prospects conclusion see the 2019 Article 17 UK Approach document.

11. Conclusions

11.1. Range	Favourable (FV)
11.2. Population	Favourable (FV)
11.3. Habitat for the species	Favourable (FV)
11.4. Future prospects	Favourable (FV)
11.5 Overall assessment of Conservation Status	Favourable (FV)
11.6 Overall trend in Conservation Status	Improving (+)
11.7 Change and reasons for change in conservation status and conservation status trend	a) Overall assessment of conservation status No change The change is mainly due to: b) Overall trend in conservation status No change The change is mainly due to:

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

11.8 Additional information

Conclusion on Range reached because: (i) the short-term trend direction in Range surface area is stable; and (ii) the current Range surface area is not less than the Favourable Reference Range.

Conclusion on Population reached because: (i) the short-term trend direction in Population size is increasing; and (ii) the current Population size is approximately equal to the Favourable Reference Population.

Conclusion on Habitat for the species reached because: (i) the area of occupied habitat is sufficiently large and (ii) the habitat quality is suitable for the long-term survival of the species; and (iii) the short-term trend in area of habitat is stable.

Conclusion on Future prospects reached because: (i) the Future prospects for Range are good; (ii) the Future prospects for Population are good; and (iii) the Future prospects for Habitat for the species are good.

Overall assessment of Conservation Status is Favourable because all of the conclusions are Favourable.

Overall trend in Conservation Status is based on the combination of the short-term trends for Range - stable, Population - increasing, and Habitat for the species - stable.

Overall assessment of Conservation Status has not changed since 2013.

Overall trend in conservation status was not reported for this species in 2013. However, from the information available the overall trend would have been improving in 2013 and so there has been no change since the last reporting round.

12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)

- a) Unit
- b) Minimum
- c) Maximum
- d) Best single value

12.2 Type of estimate

12.3 Population size inside the network Method used

12.4 Short-term trend of population size within the network Direction

12.5 Short-term trend of population size within the network Method used

12.6 Additional information

13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

Distribution Map




Figure 1: UK distribution map for S1309 - Common pipistrelle (*Pipistrellus pipistrellus*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available species records within the current reporting period. For further details see the 2019 Article 17 UK Approach document.

Range Map




Figure 2: UK range map for S1309 - Common pipistrelle (*Pipistrellus pipistrellus*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by The Mammal Society applying a range mapping tool as outlined in Matthews et al. (2018), to the 10km grid square distribution map presented in Figure 1. The alpha value for this species was 20km. For further details see the 2019 Article 17 UK Approach document.