European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Conservation status assessment for the species:

S1390 - Western rustwort (Marsupella profunda)

UNITED KINGDOM

IMPORTANT NOTE - PLEASE READ

- The information in this document represents the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- It is based on supporting information provided by the geographically-relevant Statutory Nature Conservation Bodies, which is documented separately.
- The 2019 Article 17 UK Approach document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Maps showing the distribution and range of the species are included (where available).
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the UK assessments. Further underpinning explanatory notes are available in the related country-level reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species).
- The UK-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

| NATIONAL LEVEL | | |
|---|---------------------|--|
| 1. General information | | |
| 1.1 Member State | UK | |
| 1.2 Species code | 1390 | |
| 1.3 Species scientific name | Marsupella profunda | |
| 1.4 Alternative species scientific name | | |
| 1.5 Common name (in national language) | Western rustwort | |

2. Maps

| 2.1 Sensitive species | No |
|----------------------------------|---|
| 2.2 Year or period | 2014-2015 |
| 2.3 Distribution map | Yes |
| 2.4 Distribution map Method used | Based mainly on extrapolation from a limited amount of data |
| 2.5 Additional maps | No |
| | |

| 3. Information related to Annex V Species (Art. 14) | | |
|---|---|----|
| 3.1 Is the species taken in the wild/exploited? | No | |
| 3.2 Which of the measures in Art.14 have been taken? | a) regulations regarding access to property | No |
| | b) temporary or local prohibition of the taking of specimens in the wild and exploitation | No |
| | c) regulation of the periods and/or methods of taking specimens | No |
| | d) application of hunting and fishing rules which take account of the conservation of such populations | No |
| | e) establishment of a system of licences for taking specimens or of quotas | No |
| | f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens | No |
| | g) breeding in captivity of animal species as well as artificial propagation of plant species | No |
| | h) other measures | No |

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

| b) Statistics/ quantity taken | Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period | | | | | |
|----------------------------------|---|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Season/ year 1 | Season/ year 2 | Season/ year 3 | Season/ year 4 | Season/ year 5 | Season/ year 6 |
| Min. (raw, ie. not rounded) | | | | | | |
| Max. (raw, ie. not rounded) | | | | | | |
| Unknown | No | No | No | No | No | No |

3.4. Hunting bag or quantity taken in the wild Method used

3.5. Additional information

BIOGEOGRAPHICAL LEVEL

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

Atlantic (ATL)

4.2 Sources of information

British Bryological Society, 2018. Database to the end of 2017.

Blockeel, T.L., Bosanquet, S.D.S., Hill, M.O. & Preston, C.D. 2014. Atlas of British & Irish Bryophytes. Pisces Publications, Newbury.

Porley, R.D. 2013. Marsupella profunda account in England's Rare Mosses and

Liverworts, Princeton University Press, Oxfordshire.

5. Range

5.1 Surface area (km²) 187.61 5.2 Short-term trend Period 2007-2014

5.3 Short-term trend Direction Decreasing (-)

5.4 Short-term trend Magnitude b) Maximum a) Minimum

5.5 Short-term trend Method used Based mainly on extrapolation from a limited amount of data

5.6 Long-term trend Period

5.7 Long-term trend Direction 5.8 Long-term trend Magnitude b) Maximum a) Minimum

5.9 Long-term trend Method used

5.10 Favourable reference range a) Area (km²) 1310

b) Operator c) Unknown

The FRR is the same as in 2013. The value is considered to d) Method be large enough to support a viable population and no

lower than the range estimate when the Habitats Directive came into force in the UK. For further information see the 2019 Article 17 UK Approach document.

5.11 Change and reason for change in surface area of range

Genuine change Use of different method

The change is mainly due to: Use of different method

5.12 Additional information

Short term trend in the range surface area is based on incomplete survey information. This is because not all localities where Marsupella profunda was recorded during the 2007-12 reporting period, or areas of suitable habitat, have been surveyed during the 2013-18 reporting round (in particular those areas within the working china clay quarries). Expert opinion considers that there has been some reduction in range since 2013, but the greatest difference between the current range and the range in 2013 is due to change in method and lack of survey information. The current range is more than 10% below the FRR. For further information see the 2019 Article 17 UK Approach document.

6. Population

6.1 Year or period 2007-2014

6.2 Population size (in reporting unit)

a) Unit

- number of map 1x1 km grid cells (grids1x1)
- b) Minimum
- c) Maximum
- d) Best single value

6.3 Type of estimate

Minimum

6.4 Additional population size (using population unit other than reporting unit)

- a) Unit
- b) Minimum
- c) Maximum
- d) Best single value

- 6.5 Type of estimate
- 6.6 Population size Method used

Based mainly on extrapolation from a limited amount of data

Based mainly on extrapolation from a limited amount of data

- 6.7 Short-term trend Period
- 2007-2014
- 6.8 Short-term trend Direction
- Decreasing (-)
- 6.9 Short-term trend Magnitude
- a) Minimum
- b) Maximum
- c) Confidence interval

6.10 Short-term trend Method used

- 6.11 Long-term trend Period
- 6.12 Long-term trend Direction
- 6.13 Long-term trend Magnitude
- a) Minimum
- b) Maximum
- c) Confidence interval

6.14 Long-term trend Method used

6.15 Favourable reference population (using the unit in 6.2 or 6.4)

a) Population size

21 with unit number of map 1x1 km grid cells (grids1x1)

- b) Operator
- c) Unknown
- d) Method

The FRP has changed since 2013. The value is considered to be large enough for the population to be viable and no lower than the population estimate when the Habitats Directive came into force in the UK. The UK FRP in 2013 was an operator 'more than' but it is considered that 21 1km squares (8 localities - the population in 2007) is a viable population. For further details please see the 2019 Article 17 UK Approach document and relevant country-level reporting information.

6.16 Change and reason for change in population size

Genuine change Use of different method

Use of different method The change is mainly due to:

6.17 Additional information

As not all the localities where the species was recorded in 2007-12 were surveyed in 2013-18 it is possible that the decrease in population may be less than that recorded. However, expert opinion is that some decrease has occurred. The current population is more than 25% below the FRP. For further information see the 2019 Article 17 UK Approach document.

7. Habitat for the species

7.1 Sufficiency of area and quality of occupied habitat

a) Are area and quality of occupied habitat sufficient (for long-term survival)?

No

Unknown

b) Is there a sufficiently large area of unoccupied habitat of suitable quality (for long-term survival)?

7.2 Sufficiency of area and quality of occupied habitat Method used

Based mainly on expert opinion with very limited data

7.3 Short-term trend Period

2007-2014

7.4 Short-term trend Direction

Unknown (x)

7.5 Short-term trend Method used

Based mainly on expert opinion with very limited data

7.6 Long-term trend Period

7.9 Additional information

7.7 Long-term trend Direction

7.8 Long-term trend Method used

Not all localities where the species was recorded in 2007-2012 were surveyed during this reporting round, which may have affected the observed decrease in occupied 1x1 km grids. However, suitable habitat for the species has been declining and expert opinion is that declining china clay production at active clay

pit sites may have also reduced the area of suitable habitat.

8. Main pressures and threats

8.1 Characterisation of pressures/threats

| Pressure | Ranking |
|---|--------------|
| Natural succession resulting in species composition change (other than by direct changes of agricultural or forestry practices) (LO2) | Н |
| Mining and extraction activities not referred to above (C15) | M |
| Sports, tourism and leisure activities (F07) | M |
| | |
| Threat | Ranking |
| Threat Natural succession resulting in species composition change (other than by direct changes of agricultural or forestry practices) (LO2) | Ranking H |
| Natural succession resulting in species composition change (other than by direct changes of agricultural or forestry | |

8.2 Sources of information

8.3 Additional information

Marsupella profunda occurs in disused and active china clay quarries, and is reliant on the quarrying to produce fresh areas of substrate for colonisation. A cessation of this quarrying activity would pose a threat to the species, in particular due to the vulnerability of Marsupella to being out-competed by more vigorous taller vegetation.

Marsupella profunda occurs in disused and active china clay quarries, and is reliant on the quarrying to produce fresh areas of substrate for colonisation. In some cases this quarrying activity may also adversely affect colonies of Marsupella, however this threat is distinctly less than that posed by the loss of suitable habitat caused by a cessation of quarrying activity.

Off-road motorbiking activities may cause damage to Marsupella profunda colonies in disused china clay quarries.

9. Conservation measures

9.1 Status of measures a) Are measures needed?

Yes

b) Indicate the status of measures

Measures identified and taken

9.2 Main purpose of the measures

Maintain the current range, population and/or habitat for the species

9.3 Location of the measures taken

Both inside and outside Natura 2000

9.4 Response to the measures

Medium-term results (within the next two reporting periods, 2019-2030)

9.5 List of main conservation measures

Management of habitats (others than agriculture and forest) to slow, stop or reverse natural processes (CL01)

9.6 Additional information

10. Future prospects

10.1 Future prospects of parameters

a) Range Unknown
b) Population Unknown
c) Habitat of the species Unknown

10.2 Additional information

Future trend of Range is Unknown; Future trend of Population is Unknown; and Future trend of Habitat for the species is Unknown. For further information on how future trends inform the Future Prospects conclusion see the 2019 Article 17 UK Approach document.

11. Conclusions

11.1. Range

11.2. Population

11.3. Habitat for the species

11.4. Future prospects

11.5 Overall assessment of Conservation Status

11.6 Overall trend in Conservation Status

11.7 Change and reasons for change in conservation status and conservation status trend

Unfavourable - Bad (U2)

Unfavourable - Bad (U2)

Unfavourable - Inadequate (U1)

Unknown (XX)

Unfavourable - Bad (U2)

Deteriorating (-)

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

11.8 Additional information

Conclusion on Range reached because: (i) the short-term trend direction in Range surface area is decreasing by 1% per year or less; and (ii) the current Range surface area is more than 10% below the Favourable Reference Range. Conclusion on Population reached because: (i) the short-term trend direction in Population size is decreasing by 1% per year or less; and (ii) the current Population size is more than 25% below the Favourable Reference Population. Conclusion on Habitat for the species reached because: (i) the area of occupied habitat is not sufficiently large and (ii) the habitat quality is not adequate for the long-term survival of the species; and (iii) the short-term trend in area and quality of habitat is unknown.

Conclusion on Future prospects reached because: (i) the Future prospects for Range are unknown; (ii) the Future prospects for Population are unknown; and (iii) the Future prospects for Habitat for the species are unknown.

Overall assessment of Conservation Status is Unfavourable-Bad because one or more of the conclusions are Unfavourable-Bad.

Overall trend in Conservation Status is based on the combination of the short-term trends for Range - decreasing , Population - decreasing, and Habitat for the species - unknown.

Overall assessment of Conservation Status for this species has not changed since 2013.

Overall Trend in Conservation Status has not changed since 2013.

12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)

a) Unit number of map 1x1 km grid cells (grids1x1)

- b) Minimum
- c) Maximum
- d) Best single value 0

12.2 Type of estimate

12.3 Population size inside the network Method used

Minimum

Based mainly on extrapolation from a limited amount of data

12.4 Short-term trend of population size within the network Direction

Decreasing (-)

12.5 Short-term trend of population size within the network Method used

Based mainly on extrapolation from a limited amount of data

12.6 Additional information

Due to a survey for Marsupella profunda not having occurred in the SACs in the 2013-2018 reporting round it is considered likely that the figure of '0' does not fully represent the population of the species within the SACs.

13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

Distribution Map



Figure 1: UK distribution map for S1390 - Western rustwort (*Marsupella profunda*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available species records within the current reporting period. For further details see the 2019 Article 17 UK Approach document.

Range Map



Figure 2: UK range map for S1390 - Western rustwort (*Marsupella profunda*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by applying a bespoke range mapping tool for Article 17 reporting (produced by JNCC) to the 10km grid square distribution map presented in Figure 1. The alpha value for this species was 50km. For further details see the 2019 Article 17 UK Approach document.

Explanatory Notes

| Field label | Note |
|--|--|
| 2.1 Sensitive species | A small and inconspicuous species that is not considered to be at risk from collecting, hence not sensitive. |
| 2.2 Year or Period | Species records from 2014 only have been used for this reporting round (no records of the species were available from 2013 or 2015-2018). |
| 2.3 Distribution map | 1 x 1 km grid map, produced by Natural England. |
| 2.4 Distribution map; Method used | Records of Marsupella profunda on the British Bryological Society database. |
| Species name: Marsupella pro | funda (1390) Region code: ATL |
| Field label | Note |
| 5.2 Short term trend; Period | 2007 to 2014 (no records of the species were available from 2013 or 2015-2018). |
| 5.3 Short term trend; Direction | This trend covers the years 2007 to 2014. There were fewer records in this 2013-18 reporting round, with 7 occupied 1 x 1 km squares (3 localities) recorded in 2014, compared to $211x1$ km squares (8 localities) in 2007-12. However not all the localities where Marsupella was recorded in 2007-12 were surveyed in 2013-18, and it thus possible that the decrease may be less than that recorded. |
| 5.11 Change and reason for change in surface area of range | Some observed reduction in range is likely to have been caused by not all known sites for Marsupella profunda being surveyed in 2013-2018. However expert opinion is that there has been some reduction, due to both vegetation succession on sites where there is no active quarrying for china clay, and a reduction in suitable habitat being created in active quarries. |
| 5.12 Additional information | It is considered that change in range is best assessed as Unknown as not all localities where Marsupella profunda was recorded in 2007-12 or areas of suitable habitat were surveyed during this reporting round, in particular those areas within the working china clay quarries. |
| 6.1 Year or Period | 2007 to 2014 (no records of the species were available from 2013 or 2015-2018). |
| 6.2 Population size | A total of 7 1x1 km grid squares were recorded as supporting Marsupella profunda in England in the 2013-18 reporting round, with records from 2014 only. |
| 6.7 Short term trend; Period | 2007 to 2014 (no records of the species were available from 2013 or 2015-2018). |
| 6.8 Short term trend; Direction | There were fewer records in this 2013-18 reporting round, with 7 occupied 1 x 1 km squares (3 localities) recorded in 2014, compared to 21 1 x 1 km squares (8 localities) in 2007-12. Not all the localities where Marsupella was recorded in 2007-12 were surveyed in 2013-18, and it thus possible that the decrease may be less than that recorded, however expert opinion is that some decrease has occurred. |
| 7.1 Sufficiency of area and quality of occupied habitat | Suitable habitat for the species has been declining at Lower Bostraze due to vegetation succession, and expert opinion is that declining china clay production at active clay pits sites is also likely to have reduced the area of suitable habitat. |
| 8.1 Characterisation of pressures/ threats | Marsupella profunda occurs in disused and active china clay quarries, and is reliant on quarrying to produce fresh areas of substrate for colonisation. A cessation of this quarrying activity would pose a threat to the species, in particular due to the vulnerability of Marsupella to being out-competed by more vigorous taller vegetation. |

| 9.1 Status of measures | Some conservation measures for Marsupella profunda have taken place, including the designation of SACs, however further measures are needed to conserve the species, in particular the maintenance and creation of bare china clay substrates suitable for colonisation, both within and outside the SAC areas. |
|--|---|
| 9.5 List of main conservation measures | Marsupella profunda occurs in disused and active china clay quarries, and is reliant on quarrying to produce fresh areas of substrate for colonisation. Continuation of this quarrying activity is thus important to provide suitable habitat for the species. |
| 10.2 Additional information | Future prospects have been assessed as Unknown as not all localities where Marsupella profunda was recorded in 2007-12 or areas of suitable habitat were surveyed during this reporting round, in particular those areas within the working china clay quarries. |
| 11.8 Additional information | Not all localities where Marsupella was recorded in 2007-2012 were surveyed during this reporting round, which may have affected the observed decrease in occupied 1x1 km grids. However suitable habitat for the species has been declining at Lower Bostraze due to vegetation succession, and expert opinion is that declining china clay production at active clay pits sites may have also reduced the area of suitable habitat. |
| 12.1 Population size inside the pSCIs, SCIs and SACs network | The figure of '0' is likely to be much due to a survey of SACs not having occurred within the 2013-2018 reporting round. |